

Qy 421 cggcctccccgcgcacgaagcgccgtagcagagcacttgtcacagggcggtgccctaca 480



[illegible]

Db	1761	TCCAGTATCCCATCTATATCTTTCAATAAAATGGAACCTTTTGTCTCTATGAATAGA	1820
Qy	1553	tgctcaagtcgatctactctacttttgatgtattggcctcatigccttgatcattc	1612
Db	1821	AGTCAACATCTCCTGAATATGTTCTGGTGTGTGGCTGGACCAACATAGTGAATGT	1880
Qy	1613	aataaaatgggtgtgtacacccaataaaaaa	1642
Db	1881	TATGTAGTGAAGTTAAAAAATAAAAAA	1910
RESULT	3		
ID	X09010/c		
ID	X09010 standard; cDNA; 1272 BP.		
AC	X09010;		
DT	14-JUN-1999 (first entry)		
DE	Bn-3a polynucleotide.		
KW	Bn-3a; Bcl-2; neurons; neuronal cells; apoptosis; cell death; CNS;		
KW	PNS; central nervous system; parasympathetic nervous system;		
KW	development; injury; neurotrophic factor; nerve growth factor; NGF;		
KW	ciliary neurotrophin factor; CNF; brain-derived neurotrophic factor;		
KW	BNF; neurotrophin; NT-3; NT-4; NT-5; neurodegenerative disease;		
KW	familial dysautonomia; infantile muscular dystrophy;		
KW	Parkinson's disease; Alzheimer's disease; ss.		
KW	Homo sapiens.		
FS	Key	Location/Qualifiers	
FT	CDS	1..1272	
FT		/tag= a	
FT		/product= Bn_3a_polypeptide	
PN	W090905272-A1.		
PD	04-FEB-1999.		
PF	27-JUL-1998; G02228.		
PR	10-DEC-1997; US-988476.		
PR	25-JUL-1997; GB-015823.		
PA	(UNLO ) UNIV COLLEGE LONDON.		
PI	Latchman DS, Smith MD;		
WPI	99-142928/12.		
DR	P-PSDB; W96262.		
PT	New polypeptide comprising transcription factor Bn-3a, or its		
PT	derivative - useful for treating nervous system diseases, preventing		
PT	cellular apoptosis and increasing nerve regeneration following		
PT	neuronal damage		
PS	Disclosure; Page 61-62; 68pp; English.		
CC	Over expression of transcription factor Bn-3a can protect neuronal		
CC	cells from apoptosis. Bn-3a also specifically activates expression		
CC	of the Bcl-2 gene in neuronal cells and this activation is mediated		
CC	via a Bn-3a response element in the 5' regulatory region of the		
CC	Bcl-2 gene. Both the anti-apoptotic effect of Bn-3a and its ability		
CC	to activate expression of Bcl-2 are mediated by the N-terminal		
CC	domain of Bn-3a. Members of the Bcl-2 family perform critical		
CC	roles in the regulation of selective apoptosis during development of the		
CC	nervous system. The stimulation of Bcl-2 expression by Bn-3a in a		
CC	neuron specific manner and consequent protection of neuronal cells		
CC	from apoptosis suggests that Bn-3a may co-ordinate some aspects of		
CC	neuronal reorganisation during development or following injury. The		
CC	elevation of Bn-3a expression by either pharmacological means		
CC	(compositions comprising one other therapeutic polypeptide e.g.		
CC	neurotrophic factors, nerve growth factor (NGF), ciliary neurotrophic		
CC	factor (CNTF), brain-derived neurotrophic factor (BNF), and		
CC	neurotrophins NT-3 and NT-4/5) or gene therapy may represent a		
CC	method for treating human diseases associated with excessive		
CC	neuronal cell death and/or lack of nerve regeneration, especially		
CC	neurodegenerative diseases such as familial dysautonomia and		
CC	infantile muscular dystrophy, and Parkinson's and Alzheimer's		
CC	disease.		
SQ	Sequence	1272 BP; 202 A; 468 C; 452 G; 150T;	

Query Match	3.9%	Score 64;	DB 1;	Length 1272;
Best Local Similarity	47.9%	Pred. No.	0.00049;	

Qy 47 cgtttcctcgtggcaatggtacaccagagcaacqgc-cacqqcgaqqccqccqccqcc 105















[illegible]

```

RESULT 12
Q15154 Q15154 standard; DNA; 4044 BP.
ID15154 AC Q15154;
DE 16-MAR-1992 (first entry)
DE Plasmid PHAS2-MCP containing SHV-1 major capsid protein gene.
KW P1g; recombinant vaccinia virus; PHAS2-MCP; swine herpesvirus;
KW mad itch; ss.
OS Pseudorabies virus.
FH Key Location/Qualifiers
FT cds 10..3846
FT /*tag= a
FT /note= "encodes part of SHV-1 MCP"
FT misc_feature 1..9
FT /*tag= b
FT /note= "HAS2 vector and linker"
FT misc_feature 4021..4044
FT /*tag= c
FT /note= "linker and HAS2 vector"
PN J03247285-A.
PD 05-NOV-1991.
PD PF 27-FEB-1990; 0456888.
PR PR 27-FEB-1990; JP-0456888.
PA (NORO ) NORINSHO.
PA (MITU ) MITSUBISHI KASEI CORP.
DR WPI; 91-366332/50.
DR PT Capsid protein of swine herpes virus type I - used for the
PT diagnosis, prevention and treatment of Aujeszky disease
PS Claim 3; Fig 4; 22pp; Japanese.
CC Part of the SHV-1 MCP coding sequence is inserted into a vaccinia
CC virus. It encodes amino acids 53 to 1330 of the major capsid
CC protein (see R15444 for full-length sequence).
CC See also Q15153.
SQ Sequence 4044 BP; 577 A; 1576 C; 1327 G; 564 T;

```

Query Match  
3.3%; Score 55.4; DB 1; Length 4044;

[illegible]

```

RESULT 13
T68715
ID T68715 standard; DNA; 15872 BP.
AC T68715;
DC 01-SEP-1997 (first entry)
DE Streptomyces venezuelae polyketide synthase vep ORF1.
KW polyketide synthase; polyhydroxyalkanoate monomer synthase;
KW polyhydroxybutyrate; biodegradable polymer; vep gene;
KW metabolic engineering; OS
OS Streptomyces venezuelae.
FH Key Location/Qualifiers
FT cds 20..13912
FT et /*tag= a
FT cds 14056..14136
FT ft /*tag= b
FT cds 14148..15827
FT ft /*tag= c
WO9722711-A1.
PN 26-JUN-1997.
PD 18-DEC-1996; U20119.
PF 19-DEC-1995; US-008847.
PR (MINU ) UNIV MINNESOTA.
PA Sherman DH, Williams MD, Xue Y;
PI WPI: 97-341701/31.
DR P-PSDB: W19629-30 AND W00918.
DR Expression cassettes for production of polyhydroxyalkanoate(s) -
PT provide wide range of biodegradable polymers for medical or
PT industrial use
PT Claim 54; Fig 23; 91pp; English.
PS Streptomyces venezuelae vep ORF1 (T68715) comprises the polyketide
CC synthase (PKS) gene cluster encoding a polyene of 12 carbons (see
CC also W19629-30 and W00918). It contains 5 PKS modules, with a 5'
CC loading module and a 3' end domain. Each of the sequenced modules
CC includes a keto-ACP, an acyltransferase, a dehydratase, a keto-
CC reductase and an acyl carrier protein domain. The gene cluster was
CC cloned using a heterologous hybridisation strategy from a genomic
CC DNA library. A novel expression cassette encoding the first module
CC from the vep gene cluster and module 7 from the Streptomyces ty1P
CC gene cluster has polyhydroxyalkanoate (PHA) monomer synthase
CC activity and can be used for PHA prodn. in host (esp. insect) cells
CC for use as a biodegradable polymer.

```

[illegible]



---



GenCore version 4.5  
Copyright (c) 1993 - 1998 CompuGen Ltd.

OM nucleic - nucleic search, using sw model

Run on: October 1, 1999, 13:50:36 ; Search time 1096.08 seconds  
(without alignments)  
2987.378 Million cell updates/sec

Title: US-09-026-400-1

Perfect score: 1660

Sequence: 1 attgactagctgttcattc.....aaaaaaaaaaaaaaaaaaaaa 1660

Scoring table: IDENTITY\_NUC

Searched: 2546578 seqs, 986266752 residues

Database :

EST:\*

- 1: em\_est1:\*
- 2: em\_est2:\*
- 3: em\_est3:\*
- 4: em\_est4:\*
- 5: em\_est5:\*
- 6: em\_est6:\*
- 7: em\_est7:\*
- 8: em\_est8:\*
- 9: em\_est9:\*
- 10: em\_est10:\*
- 11: em\_est11:\*
- 12: em\_est12:\*
- 13: em\_est13:\*
- 14: em\_est14:\*
- 15: em\_est15:\*
- 16: em\_est16:\*
- 17: em\_est17:\*
- 18: em\_est18:\*
- 19: em\_est19:\*
- 20: gb\_est1:\*
- 21: gb\_est2:\*
- 22: gb\_est3:\*
- 23: gb\_est4:\*
- 24: gb\_est5:\*
- 25: gb\_est6:\*
- 26: gb\_est7:\*
- 27: gb\_est8:\*
- 28: gb\_est9:\*
- 29: gb\_est10:\*
- 30: gb\_est11:\*
- 31: gb\_est12:\*
- 32: gb\_est13:\*
- 33: gb\_est14:\*
- 34: gb\_est15:\*
- 35: gb\_est16:\*
- 36: gb\_est17:\*
- 37: gb\_est18:\*
- 38: gb\_est19:\*
- 39: gb\_est20:\*
- 40: gb\_est21:\*
- 41: gb\_est22:\*
- 42: gb\_est23:\*
- 43: gb\_est24:\*
- 44: gb\_est25:\*
- 45: gb\_est26:\*
- 46: gb\_est27:\*
- 47: gb\_est28:\*
- 48: gb\_est29:\*
- 49: gb\_est30:\*
- 50: gb\_est31:\*
- 51: gb\_est32:\*
- 52: em\_est20:\*
- 53: em\_est21:\*

- 54: em\_est22:\*
- 55: em\_est23:\*
- 56: em\_est24:\*
- 57: em\_est25:\*
- 58: em\_est26:\*

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

SUMMARIES

Result No.	Score	Match	Length	DB	ID	Description
1	372.6	22.4	571	49	AI657244	AI657244 486092E01
2	301.8	18.2	594	49	AI619119	AI619119 486073F11
3	262.6	15.8	459	21	D48521	D48521 RICS14756A
4	240.2	14.5	410	20	D25143	D25143 RICS3343A R
5	177	10.7	464	49	AU056779	AU056779 AU056779
6	150	9.0	446	20	Z38084	Z38084 ATTS4224 Ve
7	136.8	8.2	470	23	T76370	T76370 11148 Lambd
8	122.8	7.4	343	20	D23043	D23043 RICC2086A R
9	117.8	7.1	352	23	T13684	T13684 1849 Lambda
10	106.6	6.4	719	40	AA980369	AA980369 ua52c12.r
11	102.4	6.2	283	49	AI621525	AI621525 486092E01
12	95.6	5.8	512	34	AA511748	AA511748 vj39f11.r
13	90.6	5.5	405	20	Z38047	Z38047 ATTS4207 Ve
14	86.8	5.2	347	23	T41757	T41757 10338 Lambd
15	85	5.1	415	48	AI597379	AI597379 vj29f11.Y
16	84.8	5.1	757	40	C91185	C91185 C91185 Dict
17	77.8	4.7	559	33	AA394723	AA394723 26506 Lam
18	77.6	4.7	536	47	AI487927	AI487927 EST246249
19	75	4.5	801	42	AI116859	AI116859 ue29d01.Y
20	74	4.5	729	41	AI048570	AI048570 ud61e01.Y
21	66.6	4.0	521	42	AI116355	AI116355 uf02b01.Y
22	63.6	3.8	528	25	N97159	N97159 22338 Lambd
23	63.4	3.8	549	26	W43288	W43288 22681 Lambd
24	60.2	3.6	664	41	AI055475	AI055475 coau0004B
25	59.8	3.6	699	45	AI386363	AI386363 mo05a03.Y
26	57.6	3.5	591	28	AA087374	AA087374 mn96b03.r
27	57.4	3.5	288	36	AA650709	AA650709 30823 Lam
28	57	3.4	437	20	T53940	T53940 yb85b03.r1
29	56	3.4	248	36	C72041	C72041 C72041 Rice
30	53.2	3.2	644	40	AA979730	AA979730 MEST1-B1.
31	53.2	3.2	618	40	AA979961	AA979961 MEST4-C10
32	53.2	3.2	621	40	AA980005	AA980005 MEST4-C5.
33	52.4	3.2	232	20	D41250	D41250 RICS3617A R
34	52.2	3.1	870	48	AI562691	AI562691 TENS2678
35	51.8	3.1	664	48	AI550776	AI550776 mn96b03.Y
36	51.6	3.1	502	36	AA395846	AA395846 25834 Lam
37	50.4	3.0	719	38	AA752790	AA752790 97AS0372
38	50.4	3.0	779	48	AI562604	AI562604 TENS2761
39	50.4	3.0	557	48	AI601030	AI601030 486096D01
40	49.2	3.0	623	31	AA292171	AA292171 zt50d08.r
41	48.8	2.9	328	42	AI138530	AI138530 gd80f01.x
42	48.8	2.9	650	46	AI442831	AI442831 sa27c08.x
43	48.8	2.9	198	49	AI626616	AI626616 fc05e10.x
44	48.6	2.9	480	42	AI132468	AI132468 ue96e10.x
45	48.4	2.9	461	49	AU056780	AU056780 AU056780

ALIGNMENTS

RESULT 1

AI657244

LOCUS

DEFINITION

AI657244

ACCESION

NID

VERSION

AI657244 571 bp mRNA EST 05-MAY-1999  
486092E01.y1 486 - leaf Primordia cdna library from Hake lab Zea  
mays CDNA, mRNA sequence.  
AI657244  
94753339  
AI657244.1 GI:4753339

Qy	737	ccaaacaatcgtgcgcagcggtttactcttaaccacattctgccaaggtcgcgcgaggtg	796
Db	478	CCCAACAACCCCTTGC	536
Qy	797	gcaaggaagctcggaatatattggtgatcgctgacga	831
Db	537	GCAAGGAAGCTTGGAATACTGGTCATGCTGATGA	571
RESULT 2			
LOCUS	AI619119/c		
DEFINITION	AI619119 594 bp mRNA EST 21-APR-1999		
ACCESSION	486073F11.x2 486 - leaf primordia cDNA library from Hake lab Zea mays cDNA, mRNA sequence.		
NID	AI619119		
VERSION	94628245		
KEYWORDS	AI619119.1 GI:4628245		
SOURCE	EST.		
ORGANISM	Zea mays.		
REFERENCE	Zea mays.		
AUTHORS	Eukaryota; Viridiplantae; Streptophyta; Embryophyta; Tracheophyta; euphyllophytes; Spermatophyta; Magnoliophyta; Liliopsida; Poales; Poaceae; Zea.		
TITLE	1 (bases 1 to 594)		
JOURNAL	Walbot,V.		
COMMENT	Maize ESTs from various cDNA libraries sequenced at Stanford University Unpublished (1999) On Mar 10, 1998 this sequence version replaced gi:2948200.  Contact: Walbot V Department of Biological Sciences Stanford University 855 California Ave, Palo Alto, CA 94304, USA Tel: 650 723 2227 Fax: 650 725 8221 Email: walbot@stanford.edu Plate: 486073 row: F column: 11.  FEATURES source 1..594 /organism="Zea mays" /cultivar="B73" /db_xref="taxon:4577" /clone_lib="486 - leaf primordia cDNA library from Hake lab" /tissue_type="leaf primordia" /dev_stage="P7-P11 leaf" /lab_host="E.coli XL1-Blue MFR/" /note="Organ: shoot; Vector: Lambda zap; Hake lab cDNA library."		
BASE COUNT	172 a 113 c 112 g 197 t		
ORIGIN			
Query Match	18.2%; Score 301.8; DB 49; Length 594;		
Best Local Similarity	78.8%; Pred. No. 6.7e-43;		
Matches	360; Conservative 0; Mismatches 97; Indels 0; Gaps 0;		
Qy	996	aaactagatctctacgtctattacgaattacccttaattgtctcaacgaccgcgaacct	1055
Db	589	AAACAAGATCATGTGCATGCATCACAAACTTCCTTAACGTTTCAACAGATCCGCAACTT	530
Qy	1056	tcggtcaggaagctcttcctaaattcttgaacacaaagcagatttcttaagagga	1115
Db	529	TTGTTTCAGGAGGCGCTTCCCAATTCCTTAAGAACACAAAGGAAGATTTTTCAGAGGA	470
Qy	1116	ttattggtctactaaaaggaatcatcagagatatgtttatagggaataaagaaaaacaaat	1175
Db	469	TCATTGCTGCTAGCGGAAACATCAGAGATATGTTTCAGCGGAATAAAGACATCAAGT	410
Qy	1176	attatagctgtctctcaacagcagaaggtatcgatgtttgttaatggtcaactaacattac	1235
Db	409	GCATCACATGCCCTCCCAAGCCAGAAGTTCCATGTTTGTCTATGGTGAACATTAATTTGT	350



Qy	1236	atctttggaggagatccatgacgacatagatttttctgcaagctcgcgaaggagaat	1295
Db	349	ATCTTTTGGAGAGCATCCATGATGATATGATTTTGTGCAAGCTGGCAAGAAGAGT	290
Qy	1296	cagtaattttatgtccgagaggtctcttggaatggaaaaattgggttcgttacttttg	1355
Db	289	CCGTGATTTTGTCTCCAGGAGAGTGTTCGGGAATGGAAAACCTGGATCCGTATCATCTTCG	230
Qy	1356	cctgctgtccatctcttcttcaagatggactcgaaagggtcgaatcattctgtcaaa	1415
Db	229	CCATGTGATTCATCTCTCTTCTTGATGGTCTTGAGAGGCTGAAATCTTCTGCCAAGGC	170
Qy	1416	acaagaagaattctataaatggtgttagttgta	1452
Db	169	ATAAGAAAGAAGATTTGCTTAATGGCCATTAACATA	133
RESULT	3		
D48521			
LOCUS	D48521	459 bp	mRNA
DEFINITION	R1CS14756A	Rice green shoot	Oryza sativa cDNA, EST
ACCESSION	D48521		
NID	g702230		
VERSION	D48521.1	GI:702230	
KEYWORDS	EST.		
SOURCE	Oryza sativa.		
ORGANISM	Oryza sativa		
REFERENCE	1 (bases 1 to 459)		
AUTHORS	Sasaki,T., Miyao,A. and Yamamoto,K.		
TITLE	Rice cDNA from callus 1995		
JOURNAL	Unpublished (1995)		
COMMENT	Contact: Takuji Sasaki National Institute of Agrobiological Resources Rice Genome Research Program 2-1-2 Kannondai, Tsukuba Ibaraki, Japan 305 Tel: 0298-38-7441 Fax: 0298-38-7468 Email: tsasaki@abr.affrc.go.jp Insert Length: 852 Std Error: 0.00 High quality sequence stop: 405.		
FEATURES	Location/Qualifiers		
source	1. .459 /organism="Oryza sativa" /strain="Nipponbare" /db_xref="taxon:4530" /clone_lib="Rice green shoot" /note="Green shoot (8 days old)"	134 a 86 c 105 g 132 t 2 others	
BASE COUNT			
ORIGIN			
Query Match	15.8%	Score 262.6; DB 21;	Length 459;
Best Local Similarity	74.4%;	Pred. No. 3.5e-36;	
Matches 343;	Conservative	0; Mismatches 116;	Indels 2; Gaps 1
Qy	672	agatcacatcgactcgctggaatccatcgcgacaagaacacacccgcgatggtcatca	731
Db	1	AGATTGACCTTAACCTCCCTAGAAATCTATTGCGGCAAGAACAACACTGCGATGATCATCA	60
Qy	732	taaacccaacaacatccgtgcgcgcgtttactcctacgacctgtgccaaggtcgcg	791
Db	61	TAAATCCCAATATCCATCCGGAATGTGTACACTTACGACATTTATCCAAAGTGGCAG	120
Qy	792	aggtgcaaggagctcggaattggtgatcgctgacgaggtttaaggcaaacgtgttc	851
Db	121	AGGTAGCAAGGAGCTTGGGNTATTGGTAATTTACTTGATGAGGTGTATGGTAATTTGGTTT	180

463	QY	acaggcgctgcctacaagctatcgccgcgacgaogtcttctcacccgccgcgcgaactca	522
Db			
61	Db	CTGTGATCTTCCTACAAAGCTTTGCACAGATGATATTTTCTCATCTCGAGGTACCCA	120
523	QY	ggcgatcgaagtcataatccgggtgctggcccgagactcgccgcgaacatactgcttc	582
Db			
121	Db	AGCAATCGAGATGTTATGCTGTTTTGGCCA---ACCAGGTGCCAATATATTGCTTCC	177
583	QY	ccggccaggctatccaaattacgaagcgcgcgagcgccattcaccaagctggaggtcccgcca	642
Db			
178	Db	AAGSCCCGGGTACCCAAACATGAGACACATGCGGTGTCCACAGGATGGAAGTCGGCT	237
643	QY	cttcgacctatcccgacaaagggtgggagatgcacatcgactcgctggaaatccatcgc	702
Db			
238	Db	CTATGATCTTGTTCACAGAGAGAGATGGAGATTAACTTTGAAGCTGTTGAAGCTTAGC	297
703	QY	cgaagaagacaccacccgagatggtcatataaaccccaacaatccgctgcgcgacggtta	762
Db			
298	Db	AGATGAGAATACTGTTGCAATAGTATTACTTAACCCCAATAACCCCTGTGTGTAATGTGA	357
763	QY	ctcctacgacctctggccaaggtcgcgagggtggcagaaggctcggaaattgggtgat	822
Db			
358	Db	CACTTATGACCATCTGCCAAGATTGCAGATACACCAAGCAAGTCCGTCGTGTAGTCAT	417
823	QY	cgctgacgaggtttaaggcaaaactggttcggcgagcccccgttt	868
418	Db	TGCTGATGAAGTATATGGTCACCTGTTTATGAAGCACTCCTTTT	463

REFERENCE	AUTHORS	TITLE	COMMENT
1 (bases 1 to 446)	Arabiopsis.		
CNRS.			
The Arabidopsis thaliana transcribed genome: the GDR cdna program			
Unpublished (1996)			
Contact: Desprez T., Anselem J., Chiappello H., Rouze P., Caboche M., Hofte H.			
INRA Versailles			
Laboratoire de Biologie Cellulaire			
Route de Saint-Cyr, 78026 Versailles Cedex France			
Email: thierry@versailles.inra.fr.			

BASE COUNT	100 a	81 c	111 g	146 t	8 others
ORIGIN					
Query Match		9.0%	Score 150;	DB 20;	Length 446;
Best Local Similarity		60.8%;	Pred. No. 7e-17;		

```

/db_xref="taxon:3702"
/map="17 cen-qter"
/clone="149B197"
/clone_lib="Lambda-PRU2"
/notes="Vector: lambda zip-Lox; Site_1: Sal; Site_2: Not;
Lambda PRU2 is a cDNA library derived from equal
quantities of 4 pools of mRNA. The mRNA sources were 1) 7
day germinated etiolated seedlings; 2) tissue culture
grown roots; 3) staged plants half with 24 hour light
cycle, half on 16 hr light, 8 hour dark- rosettes; 4)
same plants as 3 but aerial tissue (stems, flowers &
siliques). The vector is BRL's lambda zip-Lox. The cDNA
inserts were directionally cloned with Sal-Not arms using
oligo dt primed cDNA "
```

[illegible]

REFERENCE	1 (bases 1 to 343)
AUTHORS	Sasaki,T. and Minobe,Y.
TITLE	Rice CDNA from callus
JOURNAL	Unpublished (1994)
COMMENT	Contact: Takuji Sasaki National Institute of Agrobiological Resources Rice Genome Research Program 2-1-2 Kannondai, Tsukuba Ibaraki, Japan 305 Tel: 0298-38-7441 Fax: 0298-38-7468

Email: tsasaki@abr.affrc.go.jp  
PROJECT = "RGP".

# FEATURES

Location/Qualifiers  
1. 343  
/organism="Oryza sativa"  
/strain="cultivar Nipponbare, sub\_species Japonica"  
/db\_xref="taxon:4530"  
/clone="R"  
/clone\_lib="Rice callus"  
/note="Vector: pBluescript II SK+; Site 1: SalI; Site 2: NotI; cDNA prepared from rice callus mRNAs by using oligo(dT) as a primer and ligating to the SalI-NotI site of pBluescript II SK+ phagemid."  
BASE COUNT 90 a 58 c 80 g 94 t 21 others  
ORIGIN

Query Match 7.4%; Score 122.8; DB 20; Length 343;  
Best Local Similarity 70.9%; Pred. No. 3.2e-12;  
Matches 207; Conservative 0; Mismatches 81; Indels 4; Gaps 4;  
QY 780 ccaagtcgcgaggtggcaagaaagctcggaatattgtgatctgcacgaggtttacg 839  
DB 3 CCAAGTGGCAGAGGTAGCAGGAAGCTTGGGATATTGGTAATTCTGATGAGGTGTATG 62  
QY 840 gcaaacgtgtctggcagcgcgcctttatcccgatggcgctctttggcgcacattgcc 899  
DB 63 GTANTTTGGTTTGGGAGTTCNCANTTGTCCCAATNGTTACTTTGGGCACATCGTAC 122  
QY 900 cggctcttgcattggatctctgtccaaagctgtggaatgtgc-tggatggcgacttg-g 957  
DB 123 CATAANNCCNTAGGATCGCTATCAAGAGGTGGATGTCGCTAGATGGCGACTGTAG 182  
QY 958 atgggtggcggtgacagcccaagatttttagagaaacttaagatctc-tacgtcta 1016  
DB 183 NTGGNTAGCANTATGTGACCCCAAGAGACTCTACAGAAACCAAGATTGCCANCAATTAA 242  
QY 1017 ttacgaattaccattatgtctcaac-ggaccacgcaaccttgcctcaggaag 1067  
DB 243 TTAATAATTTCCTTAATGTTCAACTGNATCCAGCANCNTTTCATTACGGGAG 294

# RESULT 9

LOCUS T13684 352 bp mRNA EST 07-JAN-1998  
DEFINITION 1849 Lambda-PRL2 Arabidopsis thaliana cDNA clone 35D4T7, mRNA sequence.  
ACCESSION T13684  
NID 930780  
VERSION T13684.1 GI:930780  
KEYWORDS EST.  
SOURCE thale cress.  
ORGANISM Arabidopsis thaliana  
Eukaryota; Viridiplantae; Streptophyta; Embryophyta; Tracheophyta; euphyllophytes; Spermatophyta; Magnoliophyta; eudicotyledons; core eudicots; Rosidae; eurosids II; Brassicales; Brassicaceae; Arabidopsis.  
REFERENCE 1 (bases 1 to 352)  
AUTHORS Newman,T., deBruijn,F.J., Green,P., Keegstra,K., Kende,H., McIntosh,L., Ohlrogge,J., Raikhel,N., Somerville,S., Thomasow,M., Retzel,E. and Somerville,C.  
Genes galore: a summary of methods for accessing results from large-scale partial sequencing of anonymous Arabidopsis cDNA clones Plant Physiol. 106, 1241-1255 (1994)  
MEDLINE 95148729  
COMMENT On Nov 29, 1993 this sequence version replaced gi:430053.

Contact: Thomas Newman  
MSU-DOE Plant Research Laboratory  
Michigan State University  
MSU-DOE-PRL, Michigan State University, Plant Biology Bldg., E.  
Lansing, MI  
Tel: 517-353-0854

Fax: 517-353-9168  
Email: 22313tcn@bm.cl.msu.edu  
Seq primer: T7.

# FEATURES

Location/Qualifiers  
1. 352  
/organism="Arabidopsis thaliana"  
/strain="var columbia"  
/db\_xref="taxon:3702"  
/map="4"  
/clone="35D4T7"  
/clone\_lib="Lambda-PRL2"  
/note="Vector: lambda Zip-Lox; Site 1: Sal; Site 2: Not; Lambda PRL2 is a cDNA library derived from equal quantities of 4 pools of mRNA. The mRNA sources were 1) 7 day germinated etiolated seedlings; 2) tissue culture grown roots; 3) staged plants half with 24 hour light cycle, half on 16 hr light, 8 hour dark- rosettes; 4) same plants as 3 but aerial tissue (stems, flowers and siliques). The vector is BRU's lambda Zip-Lox. The cDNA inserts were directionally cloned with Sal-Not arms using oligo dT primed cDNA."  
BASE COUNT 77 a 61 c 83 g 118 t 13 others  
ORIGIN

Query Match 7.1%; Score 117.8; DB 23; Length 352;  
Best Local Similarity 63.1%; Pred. No. 2.3e-11;  
Matches 173; Conservative 0; Mismatches 101; Indels 0; Gaps 0;  
QY 690 tggatccatcgccgacgaagcaacaccgcgctggtcatcataaaccaacaatccgt 749  
DB 2 TCGAGGCTCTTCGACAGAAACACAGGTTCGTTGGTTGTTATAAACCCNGGTAATCCTT 61  
QY 750 cggcgagcgtttactcctcacacatctggccaaggtcgcgaggtggcaaggaagctcg 809  
DB 62 GCGGAANTCTTATAGCTACCCAGCAATTTGATGAAGATTNCGGAATCGCGAAAAAATAG 121  
QY 810 gaatttggtgatgctgacgaggtttacggcaaaactggtctcggcgagcgcccgctta 869  
DB 122 GGTTCCTTGATTNCTGATGAGGTTTACGGTCATCTTGTCTTTGGTAGCAACCGTTTG 181  
QY 870 tcccgatggcgcttcttgggcacattgcccggtcttgcattggatgctctgtccaagt 929  
DB 182 TNCCAATNGTGTGTTGGATCTATTGTNCCGTGCTTACTCTTGNTCTTTATCAAGA 241  
QY 930 cgtgatagtcctggatggcgacttgatgggt 963  
DB 242 GATGATAGTTCAGGTTTCGACCTCGGGTTNGT 275

# RESULT 10

LOCUS AA980369 719 bp mRNA EST 27-MAY-1998  
DEFINITION ua52cl2:rl Stragene mouse lung 937302 Mus musculus cDNA clone IMAGE:1350358 5' similar to gb:X52520\_cds1 TYROSINE AMINOTRANSFERASE (HUMAN);, mRNA sequence.  
ACCESSION AA980369  
NID G3158905  
VERSION AA980369.1 GI:3158905  
KEYWORDS EST.  
SOURCE house mouse.  
ORGANISM Mus musculus  
Eukaryota; Metazoa; Chordata; Craniata; Vertebrata; Mammalia; Eutheria; Rodentia; Sciurognathi; Muridae; Murinae; Mus.  
REFERENCE 1 (bases 1 to 719)  
AUTHORS Marra,M., Hillier,L., Allen,M., Bowles,M., Dietrich,N., Dubuque,T., Geisel,S., Kucaba,T., Lacy,M., Le,M., Martin,J., Morris,M., Schellenberg,K., Steptoe,M., Tan,F., Underwood,K., Moore,B., Theising,B., Wylie,T., Lennon,G., Soares,B., Wilton,R. and Waterston,R.  
The WashU-HMI Mouse EST Project  
JOURNAL Unpublished (1996)  
COMMENT On Jan 19, 1998 this sequence version replaced gi:2152585.



[illegible]

BASE COUNT	104 a	102 c	103 g	106 t
ORIGIN				

```

Query Match      5.1%; Score 85; DB 48; Length 415;
Best Local Similarity 52.1%; Pred. No. 9.7e-06;
Matches 215; Conservative 0; Mismatches 195; Indels 3; Gaps 1;

Qy 488 gccgacgactcttctccaccgcgcggaactcaggcgatcgaagtcaataatcccgggtg 547
    ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| |||
Db 4 GCTAAGGATGTCATTCTGACGAGTGGCTGCGAGTCAGGCCATTGAGCTGTGCTAGCCGTG 63

Qy 548 ctggcccaagactgcgcgcgcccaataactgcttcccggccaggctatccaaattacgag 607
    ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| |||
Db 64 TTGGCC---AATCTGGACAGAACATCTCATTCGAGGCCCGGGTTTCCCTCTACAGG 120

Qy 608 gcgagcgccattcaacaagctgaggtccggaacttcgacctcaccgccgacaagggg 667
    ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| |||
Db 121 ACATTGGCTGAGTCTATGCGGATTTGAGTCAAGCTCTACAATCTATTGCCCTGAGAAGTCT 180

Qy 668 tgggagatcgacatcgactcgtggaatccatcgccgacaagaacaccgccgcatggtc 727
    ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| |||
Db 181 TGGGAATTGATCTAAACAACCTGGAATCTCTGATCGACGAGGAAAAACAGCTTTGTCGTG 240

Qy 728 atcataaaccccaacaatccgtcgcggaagctttactctcagaccatctggccaaagtc 787
    ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| |||
Db 241 GTCAACAACCCGTCATCCCTGTGGCTCTGTGTTCAGTAAAGCGGCACCTTCAGAGAAGATT 300

Qy 788 gcgaggtggcaaggaaagctcggaattattggtgatcgctgacgaggtttacggcaaaactg 847
    ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| |||
Db 301 TTGGCAGTGGCTGAAGGCATGCGTCCCAATCTTAGCCGATGAGATCTATGGTGACATG 360

Qy 848 gtctgggagcagccccggtttatccgagtgggcgctcttttgggcacattgcccc 900
    ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| ||| |||
Db 361 GTGTTTTCAGATTGCAAAATATGAACCAATGCCCACTGCCCTCAGCACCACCAATGTCCC 413

```

Search completed: October 1, 1999, 15:03:30  
Job time: 4374 sec



GenCore version 4.5  
Copyright (c) 1993 - 1998 Compugen Ltd.

OM nucleic - nucleic search, using sw model

Run on: October 1, 1999, 13:51:37 ; Search time 1619.02 Seconds  
(without alignments)  
3260.808 Million cell updates/sec

Title: US-09-026-400-1

Perfect score: 1660

Sequence: 1 attgactagctgttcattc.....aaaaaaaaaaaaaaaaaaaaa 1660

Scoring table: IDENTITY\_NUC

Searched: 679419 seqs, 1590154680 residues

Database : GenEmbl.\*

1: gb\_ba1.\*  
2: gb\_ba2.\*  
3: gb\_om.\*  
4: gb\_ov.\*  
5: gb\_pat.\*  
6: gb\_ph.\*  
7: gb\_p11.\*  
8: gb\_p12.\*  
9: gb\_p13.\*  
10: gb\_p14.\*  
11: gb\_p15.\*  
12: gb\_p16.\*  
13: gb\_p17.\*  
14: gb\_p18.\*  
15: gb\_p19.\*  
16: gb\_p20.\*  
17: gb\_p21.\*  
18: gb\_p22.\*  
19: gb\_p23.\*  
20: gb\_p24.\*  
21: gb\_p25.\*  
22: gb\_p26.\*  
23: gb\_p27.\*  
24: gb\_p28.\*  
25: gb\_p29.\*  
26: gb\_p30.\*  
27: gb\_p31.\*  
28: gb\_p32.\*  
29: gb\_p33.\*  
30: gb\_p34.\*  
31: gb\_p35.\*  
32: gb\_p36.\*  
33: gb\_p37.\*  
34: gb\_p38.\*  
35: gb\_p39.\*  
36: gb\_p40.\*  
37: gb\_p41.\*  
38: gb\_p42.\*  
39: gb\_p43.\*  
40: gb\_p44.\*  
41: gb\_p45.\*  
42: gb\_p46.\*

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

# SUMMARIES

Result No.	Query Match	Length	ID	Description
1	144.2	8.7	2362 12 RNTATR	X02741 Rat mRNA fo

2	137.4	8.3	103495	8	ATAC006585
3	134.6	8.1	2004	12	RATTATRA
4	133.6	8.0	93695	7	ATF209
5	126.2	7.6	61384	8	ATAC007048
6	125	7.5	2754	10	HSTATR
7	125	7.5	2051	10	HSTYAT
8	118.2	7.1	73840	7	AB007644
9	110.2	6.6	39524	1	SCH10
10	110.2	6.6	70475	7	AB018112
11	107.2	6.5	25970	1	SC2H4
12	104.8	6.3	3240	1	SAAJ5198
13	104.8	6.3	3240	1	SAJ5198
14	95	5.7	39399	1	MSGB1970CS
15	94.2	5.7	119430	7	ATFYD16
16	93	5.6	9150	1	ATCY279
17	89.2	5.4	7409	1	STMVBR1
18	88.6	5.3	32258	36	CEFA2D1
19	87.8	5.3	187647	34	CEY16B4
20	73.6	4.4	1663	1	TAX99521
21	67.2	4.0	1777	1	THASPC
22	67	4.0	2977	1	AF025665
23	67	4.0	7235	1	SGNUSG
24	65.6	4.0	5732	1	STMUTA
25	65.4	3.9	3824	9	HUMOCF1A
26	64	3.9	1149	9	HSBN3A2
27	63.6	3.8	38532	1	SCE94
28	63.6	3.8	3492	10	HSRDLMR
29	63.6	3.8	3492	14	G28612
30	63.4	3.8	2249	1	SPSNBCDE
31	63.4	3.8	2249	1	SPSNBCGEN
32	63.4	3.8	2028	8	SSI132828
33	62.8	3.8	17533	1	D90860
34	62.8	3.8	10663	2	AE000318
35	62.6	3.8	37445	1	SC8A6
36	62	3.7	53784	1	AMM223012
37	62	3.7	76199	2	AF040570
38	62	3.7	53789	5	A69720
39	61.6	3.7	2767	7	SSI224970
40	61.4	3.7	22789	1	SC2A11
41	61	3.7	67200	1	MTV017
42	61	3.7	3379	3	CSCLOXA
43	60.2	3.6	2984	36	TRBANTA
44	59.6	3.6	1166	1	STMNUSG
45	59.6	3.6	1506	4	CHKMARKS

## ALIGNMENTS

RESULT 1

LOCUS

RNTATR

DEFINITION

Rat mRNA for tyrosine aminotransferase (L-tyrosine: 2-oxoglutarate

amino transferase EC 2.6.1.5).

X02741 X12522 X15375

ACCESSION

NID

VERSION

KEYWORDS

SOURCE

ORGANISM

REFERENCE

AUTHORS

TITLE

JOURNAL

MEDLINE

REFERENCE

AUTHORS

TITLE

RNTATR

2362 bp

mRNA

12-SEP-1993

Rat mRNA for tyrosine aminotransferase (L-tyrosine: 2-oxoglutarate

amino transferase EC 2.6.1.5).

X02741 X12522 X15375

ACCESSION

NID

VERSION

KEYWORDS

SOURCE

ORGANISM

REFERENCE

AUTHORS

TITLE

JOURNAL

MEDLINE

REFERENCE

AUTHORS

TITLE

AC006585 Arabidops  
M18340 Rat tyrosin  
AL021749 Arabidops  
X007048 Arabidops  
X52520 Human mRNA  
X55675 H.sapiens m  
AB007644 Arabidops  
AL049754 Streptomy  
AB018112 Arabidops  
AL031514 Streptomy  
AJ005198 Streptomy  
AJ005617 Streptomy  
L78815 Mycobacteri  
X035394 Arabidops  
Z97991 Mycobacteri  
D50624 Streptomyce  
Z81081 Caenorhabdi  
AL021148 Caenorhab  
X95221 T.aquaticus  
D38459 Thermus the  
AF025665 Thermus a  
X72787 S.griseus n  
L10064 Streptomyce  
L20433 Streptomyce  
U10063 Human POU d  
AL049628 Streptomy  
X64624 H.sapiens m  
G28612 human STS S  
Y11548 S.pristinae  
X98690 S.pristinae  
AJ332828 Spermatoz  
D90860 E.coll geno  
AE000318 Escherich  
AL031013 Streptomy  
AJ223012 Amycolato  
AF040570 Amycolato  
A69720 Sequence 3  
AJ224970 Spermatoz  
AL031184 Streptomy  
X021897 Mycobacte  
X69017 Canis sp. m  
L00673 Trypanosoma  
D17465 S.coelicolo  
M31650 Chicken myr







Db	994	CATTTTGGCAA---TGAGATTCCGAGACGGGCTGCTGAAACTGAGTCACGGGATCCTGGG	1050
Qy	1045	cccagcaaccttcgttcaggaagctcttcctaaaattcttgagaacacacaaagcagattt	1104
Db	1051	ACCATGCACCATAGTCAGGGTGCTCTGAAAGAGCATCTTCAGCGAACCCCTCAGGAGTT	1110
Qy	1105	ctttaagaggattattggtctactactaaaaggaatcatcagagatatgttatagggaataaa	1164
Db	1111	CTATCAGCACACGTTAAGCTTCCTCAAGTCCAATCGGACCTCTGCTATGGGCACTGGC	1170
Qy	1165	ggaacacaaatattacgtctcctcaaacgcccagaagatcgatgttgttaagtgtcaa	1224
Db	1171	TGACCTCCTCGACTCCAG---CCGGTCGGCCCTCTCGAGCCATGTACCTATGTTGGG	1227
Qy	1225	actaaattacatcttttggaggagatccatgacgacacatagattt	1269
Db	1228	AATTGAGATGAGCATTTCCGGAAATTCGAGACGACGTGGAGTT	1272
RESULT	4		
ATF2009			
LOCUS	ATF2009	93695 bp	DNA
DEFINITION	Arabidopsis thaliana DNA chromosome 4, BAC clone F2009 (ESSAII Project).	PLN	27-OCT-1998
ACCESSION	AL021749		
NID	92842474		
VERSION	AL021749.1	GI:2842474	
KEYWORDS			
SOURCE	thale cress.		
ORGANISM	Arabidopsis thaliana		
REFERENCE	Eukaryota; Viridiplantae; Streptophyta; Embryophyta; Tracheophyta; euphyllophytes; Spermatophyta; Magnoliophyta; eudicotyledons; Rosidae; Caprales; Brassicaceae; Arabidopsis.		
AUTHORS	1 (bases 1 to 93695) Bevan, M., Koetser, P., Hempel, S., Entian, K.-D., Hoheisel, J., Mewes, H.W., Mayer, K.F.X. and Schueller, C.		
JOURNAL	Unpublished		
REFERENCE	2 (bases 1 to 93695)		
AUTHORS	EU Arabidopsis sequencing, project.		
TITLE	Direct Submission		
JOURNAL	Submitted (16-OCT-1998) MIPS, at the Max-Planck-Institut fuer Biochemie, Am Klopferspitz 18a, D-82152 Martinsried, FRG, Project Coordinator: Mike Bevan, Molecular Genetics Department, Cambridge Laboratory, John Innes Centre, Colney Lane, NR4 7UJ Norwich, UK, E-mail: michael.bevan@bbsrc.ac.uk		
FEATURES	Location/Qualifiers		
source	1..93695		
gene	/organism="Arabidopsis thaliana"		
exon	/variety="Columbia"		
	/db_xref="taxon:3702"		
	/chromosome="4"		
	18..467		
	/gene="F2009.10"		
	complement(18..467)		
	/gene="F2009.10"		
	/number=1		
CDS	complement(18..467)		
	/gene="F2009.10"		
	/codon_start=1		
	/product="hypothetical protein"		
	/protein_id="CAAL6872.1"		
	/db_xref="PID:e1250050"		
	/db_xref="PID:g2842475"		
	/db_xref="GI:2842475"		
	/db_xref="SPREMBL:O49442"		
	/translation="MTSYKLFLVIRKRRTRWACFLVYAILSSWSNLSINVTNYG ENVHSGGPAIYASVLLGAVFGVLSMAALFAIVPAIVIVISVVVTFAGAKSRK VITEGRKVEIAGAGFVRLKGFNEFALLCALLAYFVFENSYSSSS"		
exon	complement(2196..2578)		
	/gene="F2009.20"		
	/number=1		
gene	2196..4222		
	/gene="F2009.20"		

/db\_xref="SPTREMBL:O49444"  
/translation="MGLVWREDLYLMFVLMGLGFTISNGKGYVGGKDGWVPSPSED  
YHWSHRNRFVNDTLHFYAKGKDSVLEVTQYNTNTTPTLSLSLSDGSLFLSH  
SGSVEFTLSGNSCLGOKLAVKVLSTVHSHSPRHTSPSPVHQELSSPGSPGVE  
PSSNSRVPAGFATAPNSAGLVGPMVLVNSIIPIDSFLLKSLPLVVDYVAGI  
STROLQASEPVSPLSSPALLSGKEEKIIPKROKVOALKSIQSPKKVNLVA  
ALVRGMRYEDALIOQVTVKRAAQTVYVHAARANATHNKGDPDRLLVAEAFVGK  
LFGKKVYHAKRGSIISIPRCRLTVIVRETTPEEAEIARLKHNFKKSKRERQLV  
PKLIETSPIWNRGTKANRSSELVLTILDVTCVGNMEKNRLDNLNINNIYHNH  
EETCSFRNGSERIDVDADVVVDVVELLVSENIFFNGEDSAIGVTYIGEVEAEMLR  
"

intron	complement(4667. .6566) /gene="F2009.30" /number=1
intron	4942. .4986 /gene="F2009.40" /number=1
exon	4987. .5044 /gene="F2009.40" /number=2
intron	5045. .5095 /gene="F2009.40" /number=2
exon	5096. .5188 /gene="F2009.40" /number=3
exon	complement(6567. .6669) /gene="F2009.30" /number=2
intron	complement(6670. .6877) /gene="F2009.30" /number=2
exon	complement(6878. .7149) /gene="F2009.30" /number=3
intron	complement(7150. .7304) /gene="F2009.30" /number=3
exon	complement(7305. .7371) /gene="F2009.30" /number=4
intron	complement(7372. .7536) /gene="F2009.30" /number=4
exon	complement(7537. .7647) /gene="F2009.30" /number=5
intron	complement(7648. .7762) /gene="F2009.30" /number=5
exon	complement(7763. .7915) /gene="F2009.30" /number=6
intron	complement(7916. .9900) /gene="F2009.30" /number=6
exon	complement(9901. .9975) /gene="F2009.30" /number=7
intron	complement(9976. .10604) /gene="F2009.30" /number=7
tRNA	10056. .10127 /gene="F2009.30" /note="Lys-tRNA (TTT)"
exon	complement(10605. .10996) /gene="F2009.30" /number=8
intron	complement(10997. .11073) /gene="F2009.30" /number=8
exon	complement(11074. .11254) /gene="F2009.30"

gene	/number=9 13176. .17409
Query Match	8.0%; Score 133.6; DB 7; Length 93695;
Best Local Similarity	63.5%; Pred. No. 4.1e-11;
Matches 221: Conservative	0; Mismatches 124; Indels 3; Gaps 1;
Qy 440	agcgccgtagcagacgacctgtcacagggcgctgcctcacagctatcgccgcgacgacgc 499 
Db 33691	AGGCGGGTCGCAGATTACGTTGAACCGAGACTTGACGAACAGGTAAGCCCTAATGATGTA 33750
Qy 500	ttctcaccgcgcgcgaactcaggcgatcgaaagtcataatccccgggtgctggccagact 559 
Db 33751	TTCATAACTGTCGGATGCAACCAAGGGATAGTAGTTCTTCAGTCACTGGCTCGACCA 33810
Qy 560	gccggcgccaaacatactgtctcccgccaggctaccaaattacgagcgcgagcggca 619 
Db 33811	AA---CGCTAAATCCTTCTCCCAAGGCCTAGTTACCTCTACTAGGAGCTCGTCCGCTC 33867
Qy 620	ttcaacaagctggaggtccggcacttcgacatcctccccgacaagggtgggagatcgac 679 
Db 33868	TACAGTGGACTCGAGTCCGCAAGTTCGATCTCCTTCCAGAGAAAGTAATGGAGATTGAT 33927
Qy 680	atcgactcgctggaatccatcgccgacaaagaacacacccacccgagtggtcatataaccca 739 
Db 33928	CTCCAGGCATCGAAGCCATGGCAGATGAGAACACAGTTGCAATGGTTATCATAAACCCCT 33987
Qy 740	aacaatcgctgcgcgagcggtttactcctacgacctctggccaaggtc 787 
Db 33988	AACAACCCCTGTGGAATGTTTACTCTTACGATCATCTTAAGAGGTC 34035
RESULT 5	
ATAC007048	ATAC007048 61384 bp DNA PLN 25-MAR-1999
LOCUS	Arabidopsis thaliana chromosome II BAC F23N11 genomic sequence,
DEFINITION	complete sequence.
ACCESSION	AC007048
NID	94512646
VERSION	AC007048.3 GI:4512646
KEYWORDS	HTG.
SOURCE	thale cress.
ORGANISM	Arabidopsis thaliana
REFERENCE	1 (bases 1 to 61384) Lin,X., Kaul,S., Shea,T.P., Fujii,C.Y., Shen,M., VanKken,S.E., Barnstead,M.E., Mason,T.M., Bowman,C.L., Ronning,C.M., Benito,M., Carrera,A.J., Creasy,T.H., Buell,C.R., Town,C.D., Nierman,W.C., Fraser,C.M. and Venter,J.C. Arabidopsis thaliana chromosome II BAC F23N11 genomic sequence
AUTHORS	Unpublished
TITLE	2 (bases 1 to 61384) Lin,X. and Kaul,S.
JOURNAL	Direct Submission
REFERENCE	Submitted (09-MAR-1999) The Institute for Genomic Research, 9712 Medical Center Dr., Rockville, MD 20850, USA, xlin@tigr.org
AUTHORS	3 (bases 1 to 61384) Lin,X.
TITLE	Direct Submission
JOURNAL	Submitted (25-MAR-1999) The Institute for Genomic Research, 9712 Medical Center Dr., Rockville, MD 20850, USA
COMMENT	On Mar 25, 1999 this sequence version replaced gi.4389529. Address all correspondence to: Xiaoying Lin The Institute for Genomic Research 9712 Medical Center Dr. Rockville, MD 20850, USA e-mail: xlin@tigr.org BAC clone F23N11 is from Arabidopsis chromosome II and is near the molecular marker mil48.

The orientation of the sequence is from SP6 to T7 end of the BAC clone.

Genes were identified by a combination of three methods: Gene prediction programs including GRAIL (available by anonymous ftp from arthur.epm.orn.gov), Genefinder (Phil Green, University of Washington), GenScan (Chris Burge, <http://gnomic.stanford.edu/~chris/GENSCANW.html>), and NetPlantGene (<http://www.cbs.dtu.dk/netgene/cbsnetgene.html>), searches of the complete sequence against a peptide database and the Arabidopsis EST database at TIGR (<http://www.tigr.org/tdb/at.html>).

Annotated genes are named to indicate the level of evidence for their annotation. Genes with similarity to other proteins are named after the database hits. Genes without significant peptide similarity but with EST similarity are named as 'unknown' proteins. Genes without protein or EST similarity, that are predicted by more than two gene prediction programs over most of their length are annotated as 'hypothetical' proteins. Genes encoding tRNAs are predicted by tRNAscan-SE (Sean Eddy, <http://genome.wustl.edu/eddy/tRNAscan-SE/>). Simple repeats are identified by RepeatMasker (Arian Smit, <http://ftp.genome.washington.edu/RM/RepeatMasker.html>). Regions of genomic sequence that are not annotated as genes but have predicted exons by GRAIL are annotated as misc features.

## FEATURES

## source

```

Location/Qualifiers
1..61384
/organism="Arabidopsis thaliana"
/cultivar="Columbia"
/db_xref="taxon:3702"
/chromosome="II"
/map="ml148"
/clone="F23N11"
1
/feature="16680 bases before this point were not included in
the submitted sequence, but they appear in the overlapping
BAC."
1..13161
/feature="overlap with BAC clone F5H14
(AC006234:116508..129667)."
complement(69..103)
/rpt_family="AT_rich"
384..409
/rpt_family="AT_rich"
697..721
/rpt_family="AT_rich"
1012..1035
/rpt_family="AT_rich"
1423..1572
/feature="exon predicted by xgrail, quality excellent"
complement(1528..1620)
/feature="exon predicted by xgrail, quality
marginal_shadowexon"
join(<1719..1934,2049..2277,2589..>3028)
/feature="F23N11.1"
<1719..>3028
/feature="F23N11.1"
/feature="F5H14.36"
join(1719..1934,2049..2277,2589..3028)
/feature="F23N11.1"
/feature="unknown protein"
/codon_start=1
/protein_id="AAD21702.1"
/db_xref="PID:g4512647"
/db_xref="GI:4512647"
/translation="MDRQNSDDIMRFDGMASDDVLFGFDEGNOSPEFSVNLNAG
EDDGEDDDNNNSDNEKAFWQEHQQLQGLYRSTETKIRQATKALKQVKSGLY
CVCRRPVGGRSLRGEISHLRDLVAGYDCVISKSKWQCDIPAGEHFEIVDRS
GSKSEMRVTELSFAFEIATKGESEYKRLISRLPEYVGTKRLSLRILKILCIAGK
KCLRDKKHMAPWKRYMQAKLWGLTCDDRSSLSLEASVSEAMEPENWVPVAKPRVSMNLN
YDGLLGFSAGPATVAVV"
complement(2296..2324)
/rpt_family="AT_rich"
complement(2565..2585)
/rpt_family="AT_rich"

```

```

misc_feature
/feature="exon predicted by xgrail, quality
marginal_shadowexon"
/rpt_family="POLY_A"
3837..3973
/feature="exon predicted by xgrail, quality marginal"
complement(4252..4308)
/rpt_family="(TA)n"
4307..4352
/feature="exon predicted by xgrail, quality good"
complement(4870..4893)
/rpt_family="AT_rich"
5283..5408
/rpt_family="POLY_A"
complement(<5419..>5724)
/feature="F23N11.2"
complement(<5419..>5724)
/feature="F23N11.2"
/feature="predicted by genscan"
complement(5419..5724)
/feature="F23N11.2"
/feature="hypothetical protein"
/codon_start=1
/protein_id="AAD21703.1"
/db_xref="PID:g4512648"
/db_xref="GI:4512648"
/translation="MKLLIFAVIISVVFVLVSSPTKCDLQSGKINGEKEIMNM
RLGLDVSSRRILQASRYISYEALKNLPDRNRGPDQDNPYRRSCDVHSHCYRFTN"
5852..5890
/feature="exon predicted by xgrail, quality
marginal_shadowexon"
6449..6487
/rpt_family="AT_rich"
6837..6880
/rpt_family="AT_rich"
7043..7110
/rpt_family="POLY_A"
7231..7292
/rpt_family="AT_rich"
7374..7406
/rpt_family="AT_rich"
complement(7723..7789)
/rpt_family="AT_rich"
complement(7930..8000)
/feature="exon predicted by xgrail, quality good_shadowexon"
8174..8302
/feature="exon predicted by xgrail, quality excellent"
complement(8465..8586)
/feature="exon predicted by xgrail, quality good_shadowexon"
8544..8629
/feature="exon predicted by xgrail, quality good"
8702..8829
/feature="exon predicted by xgrail, quality good"
complement(8934..9057)
/feature="exon predicted by xgrail, quality good_shadowexon"
8936..9051
/feature="exon predicted by xgrail, quality excellent"
9147..9243
/feature="exon predicted by xgrail, quality excellent"
9290..9338
/feature="exon predicted by xgrail, quality marginal"
complement(9317..9461)
/feature="exon predicted by xgrail, quality good_shadowexon"
9492..9555
/feature="exon predicted by xgrail, quality marginal"
complement(9654..9780)
/feature="exon predicted by xgrail, quality
marginal_shadowexon"
9666..9814
/feature="exon predicted by xgrail, quality excellent"
complement(9912..9988)
/feature="exon predicted by xgrail, quality

```

[illegible]

RESULT	6	
HSSTATR		
LOCUS		
DEFINITION		
ACCESSION		
NID		
VERSION		
KEYWORDS		
SOURCE		
ORGANISM		
REFERENCE		
AUTHORS		
TITLE		
JOURNAL		
REMARK		
REFERENCE		
AUTHORS		
TITLE		
JOURNAL		
MEDLINE		
COMMENT		
FEATURES		
source		

CDS	/organism="Homo sapiens" /db_xref="taxon:9606" /chromosome="16q22.1" /tissue_type="placenta (genomic) and liver (CDNA)" /clone_lib="genomic (phage) + cDNA" /clone="lambda-HTA11, phcTAT2-16 + phcTAT3a-6" 97..1461 /note="tyrosine aminotransferase (AA 1-454)" /codon_start=1 /protein_id="CAA36750.1" /db_xref="PID:g36713" /db_xref="GI:36713" /db_xref="SWISS-PROT:P17735" /translation="MDPYMIQSSKNLPSILDVHNVGGRSVPCKMKGRKARWSVR PDMCAKFNPIRAIVDNMKYFNPKNMTISIGDPTFGNLPITPEVTQANKDALD SKGYAPSIIFGLTSREIRAYHYGCEAPLEAKDIYLSGCQSQAIDLCIAVLANPGQ NIYLPPGFSLLTKAESMGIEVKLYLLPEKSWEIDLKOLEYIDIKETACLIYNPS NPCGSVFLSRHLQILAVAROCVIPLADEIYGMVFSDCKVEPLATLSTDVPILSCG GIAGLRWPKRWLGWLIIHDRDIFGNRIEIDGLVKLSORILGPCTTVOGALKSLICRT PGEFYNTLSFLSKSNADLCYGALAAIPGLRPVPRSGAMLYMVGMEHPFEFENDVEF TERLVAEQSVCHLCATCFEPNFIRVIIVPEVMLEASRSIOEFCEQHVCHEAGSQSE ECKD"
misc_feature	2070..2075 /note="minor mRNA polyadenylation signal"
polyA_site	2090 /note="minor mRNA polyadenylation site"
repeat_region	2116..2130 /note="direct repeat flanking Alu element" complement(2131..2479) /note="Alu element"
misc_feature	2480..2494 /note="direct repeat flanking Alu element"
repeat_region	2738..2743 /note="major mRNA polyadenylation signal"
polyA_site	2754 /note="major mRNA polyadenylation site"
BASE COUNT	689 a 646 c 635 g 784 t
ORIGIN	
Query Match	7.5%; Score 125; DB 10; Length 2754;
Best Local Similarity	49.0%; Pred. No. 1.le-09;
Matches	422; Conservative 0; Mismatches 430; Indels 9; Gaps 3;
Qy	297 ccgtgcgctggccagcgtagacccttcggtttcccgcgcttcgcagcgccgtcg 356 
Db	311 CCATGATTCCCTGTCCATTGGGACCCTACTGTGTTTGGAACCTGCCACACCCCTG 370 
Qy	357 aggccgaagcacgcgtgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcg 416 
Db	371 AAGTTACCCAGGCAATGAAAGATGCCCTGGACTCGGCAATAATTAATGGGTATGCCCAT 430 
Qy	417 gcgtcgcgcctccccgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcct 476 
Db	431 CCATCGGCTTCCTATCCAGTCGGGAGAGATTGCTTCTTATACCACCTGCTCAGGCAC 490 
Qy	477 acaagtatcggc 536 
Db	491 CC---CTAGAAGCTAAGAGAGCTCATTTGACAAGTGCTGCAGCAAGCTATTGACCTTT 547 
Qy	537 taatcccggtcgtggccagacatgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgc 596 
Db	548 GTTTAGCTGTGTGGCCCAACCCA---GGGAGAACATCTCTGGTTTCCAAGACCTGGTTCT 604 
Qy	597 caaatcacgaggc 656 
Db	605 CTCCTACAGACTCTTGGCTGAGTCTATTGGAAATTGAGGTCAAACCTCTACAAATTGTTGC 664 
Qy	657 ccgacaagggtggagaatcgacatcgactcgcgtggaatcccatcgcgacaagaacacca 716 
Db	665 CAGAGAAATCTTGGGAAATTCAGCTGAAACCAACTGGAATATCTAATTTGTAAGAACAG 724 
Qy	717 ccgcgatggtcatataaacccaaacaatccgtgcgcgcgcgcgcgcgcgcgcgcgcgcgc 776 



```

Db 725 CTTGTCTCATTTGCAATAATCCATAAACCCTGTGGGTGAGTGTTCAGCAAAACGTCATC 784
QY 777 tggccaagtcgaggggaggaagctcggaatattgggtgatcgctgacagaggttt 836
Db 785 TTCAGAGATTCGGCAGTGGCTGCACGGCAGTGTGCCCATCTTAGCTGTGATGAGATCT 844
QY 837 acggcgaactggcttgggcagcgcccgctttatcccgatggcgctctttgggcacattg 896
Db 845 ATGAGACATGGTGTTCGATTGCAATATGAACCACTGGCCACCCTCAGCACCGATG 904
QY 897 cccggcttgcattgatctctcgaagtcgtagtagtgcctggatggcgacttg 956
Db 905 TCCCCATCTCTCTGTGGAGGCTGCCAAGCGCTGGCTGTCTGGCTGGAGGTGG 964
QY 957 gatgggtggcggtgacgacccacacaaagattttagagaaactaagatctctactcta 1016
Db 965 GCTGGATCTCATATGACCGAGAGACATTTTGGCA--TGAGATCCGAGATGGGC 1021
QY 1017 ttacgaattaccettaatgtctcaacggaccacgaaccttcgttcaggagctcttcccta 1076
Db 1022 TGGTGAAGCTGAGTCAGCGCATTTTGGACCCCTGTACCATTTGCCAGGGAGCTCTGAAA 1081
QY 1077 aaattcttgagacacaaagagatttcttaaggaggattattggtctactaaagaaat 1136
Db 1082 GCATCTATGTGCGACCCCGGAGAGTTTACACAACTCTGAGCTTCTCAAGTCCA 1141
QY 1137 catcagagatatgtatagg 1157
Db 1142 ATGCTGATCTGTTATGGG 1162

RESULT 7
HSTYAT Locus HSTYAT 2051 bp mRNA PRI 13-JUL-1995
DEFINITION H sapiens mRNA for tyrosine aminotransferase.
ACCESSION X55675
NID 937501
VERSION X55675.1 GI:37501
KEYWORDS tyrosine aminotransferase.
SOURCE human.
ORGANISM Homo sapiens
Eukaryota; Metazoa; Chordata; Vertebrata; Mammalia; Eutheria;
Primates; Catarrhini; Hominiidae; Homo.
REFERENCE 1 (bases 1 to 2051)
AUTHORS Labrie,F.
TITLE Direct Submission
JOURNAL Submitted (26-SEP-1990) Labrie F., CHUL Research Centre, Laboratory
of Molecular Endocrinology, 2705 Blvd Laurier, Sainte-Foy Quebec
GLV 4G2, Canada
2 (bases 1 to 2051)
AUTHORS Serailini,G.E., Luu-The,V. and Labrie,F.
TITLE Cloning and expression of human tyrosine aminotransferase cDNA
JOURNAL Biochim. Biophys. Acta 1260 (1), 97-101 (1995)
MEDLINE 95092801
FEATURES Location/Qualifiers
1..2051
/organism="Homo sapiens"
/db_xref="taxon:9606"
/chromosome="16"
/dev_stage="adult"
/tissue_type="liver"
/clone_lib="# HL100lb. Lot # 2102"
/clone="B1"
/map="16q 22.1"
51..1415
/EC_number="2.6.1.5"
/codon_start=1
/product="tyrosine aminotransferase"
/protein_id="CAA39210.1"
/db_xref="PID:937502"
/db_xref="GI:37502"
/db_xref="SWISS-PROT:P17735"
CDS

```

```

/translation="MDPYMIOMSSKGNLPSILDVHNVYGRSSVPGKMKGRKARNSVR
PSDMAKTFNPPIRAIVDMKYAPFNKNTMISLISGIDPVTGFLNLPDPEVTOAMKDALD
SKYNGYAPSGISLRSBEIAYIHCPEAPLEAKDVLITSCQSDIDCLAVLANPGQ
NLTVPFGFSLYKTLAESMGIEVKLYNLPEKSWIDLKQLEYLIDETACLIIVNPS
NPGSVFSKRHLQKTLAARQCVPILADEIYGDWVFSCKVEPLATLSTDPILSCG
GLAKRWLPVGRWLGLIHDRDIFGNEDRLGLVLSORILGPICTIVOGALAKIILCRT
PGEFVHTLSFLKSNADLCYGAALPGLRPVPSGAMVLMVGMIEHMFPEENDEVEF
TERLVAEQSVHCLPATCFEYPNFRIVITVPVMMLEACSLQIEFCEQHYHCAESQSE
BCDK"
polyA_signal 2021..2026
BASE COUNT 531 a 493 c 496 g 531 t
ORIGIN
Query Match 7.5%; Score 125; DB 10; Length 2051;
Best Local Similarity 49.0%; Pred. No. 1.1e-09;
Matches 422; Conservative 0; Mismatches 430; Indels 9; Gaps 3;
QY 297 ccgtgtgctgcgtggccacggtgacccgtccggtgttcccgccctccgcagggccgtcg 356
Db 265 CCATGATTTCCCTGTCCATTTGGGACCCCTACTGTTTGGAAACCTGCTACAGACCCCTG 324
QY 357 aggcgaagacgcgcgtgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgc 416
Db 325 AAGTTACCCAGGCAATGAAAGATGCCCTGGACTCGGGCAATATATATGCTATGCCCAT 384
QY 417 gcgtgcgcctcccgccgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgc 476
Db 385 CCATCGGCTTCTATCCAGTGGGAGGAGATTGCTTTATTACCACCTGCTCTGAGGCAC 444
QY 477 caaagctatcgccgacgacgtctctcaccgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgc 536
Db 445 CC--CTAGAAGCTAAGGACGTCATCTGACAAGTGGCTGCGACCAAGTATTGACCTTT 501
QY 537 taatcccggtgtgccccagactgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgc 596
Db 502 GTTTAGCTGTGTGGCCAAACCA--GGCAAAACATCCTGTTTCCAGACCTGGTTTCT 558
QY 597 caaattagagcgagcgagcgacattcaacaagctgagggctccgacgttcgacctcatcc 556
Db 559 CTCTCTACAAGACTCTGGCTGAGTCTATGGGAATTGAGGTCAAACCTCTACAAATTTGTTGC 618
QY 657 ccgacaaggggtgggagatcgacatcgactcgtcgtgaatccatcgccgacaagaacacca 716
Db 619 CAGAGAAATCTTGGAAATTTGACCTGAACACTGGAATATCTAATTGATGAAGACAG 678
QY 717 ccgcatggttcataataaccacaacatccgtgcgcgcgcgcgcgcgcgcgcgcgcgcgcgcgc 776
Db 679 CTTGTCTCATTTCAATAATCCATCAACCCCTGTGGGTGAGTGTTCAGCAAAACGTCATC 738
QY 777 tggccaagtcgaggtggcgaagagctcggaatattgggtgatcgctgacagaggttt 836
Db 739 TTCAGAGATTTCTGGCAGTGGCTGCACGGCAGTGTGCCCATCTTAGCTGTGATGAGATCT 798
QY 837 acggcgaactggcttgggcagcgcccgctttatccgatggcgctctttgggcacattg 896
Db 799 ATGAGACATGGTGTTCGATTGCAATATGAACCACTGGCCACCCTCAGCACCGATG 858
QY 897 ccccggttcttccattggtatctctgtccaaagctgtgtagtgcctggatggcgacttg 956
Db 859 TCCCCATCTCTCTGTGGAGGCTGGCCAGCGCTGGCTGGTGTCTGGCTGGAGGTGG 918
QY 957 gatgggtggcggtgtacacccccacaaagatttttagagaaactaagatctctactcta 1016
Db 919 GCTGGATCTCATTCATGACCGAGAGACATTTTGGCAA---TGAGATCCGAGATGGGC 975
QY 1017 ttacgaattaccettaatgtctcaacggaccacgaaccttcgttcaggagctcttcccta 1076
Db 976 TGGTGAAGCTGAGTCAGCGCATTTTGGACCCCTGTACCATTTGTCAGGGAGCTCTGAAA 1035
QY 1077 aaattcttgagacacacaaagagatttcttaaggaggattattggtctactaaagaaat 1136
Db 1036 GCATCTATGTGCGACCCCGGAGAGTTTACCAACACTCTGAGCTTCTCAAGTCCA 1095

```



Cosmid H10 lies between 66r3 and H44 on the Aseri-H genomic restriction fragment.

## FEATURES

```

source      Location/Qualifiers
            /organism="Streptomyces coelicolor"
            /strain="A3(2)"
            /db_xref="taxon:1902"
            /clone="cosmid H10"
            1..360
            /gene="SCH10.01"
            misc_feature 1..44
            /gene="SCH10.01"
            /note="overlap with S.coelicolor cosmid St66r3 from 35058
            35101"
            <1..360
            /gene="SCH10.01"
            /note="SCH10.01, hypothetical protein, partial CDS, len:
            >119 aa; probable CDS suggested by positional base
            preference, GC frame analysis and amino acid composition"
            /codon_start=1
            /transl_table=11
            /label="SCH10.01"
            /product="hypothetical protein"
            /protein_id="CAB42010.1"
            /db_xref="PID:e1428991"
            /db_xref="PID:e1428992"
            /db_xref="PID:e1428993"
            /db_xref="GI:4753847"
            /translation="DPRIAAALAGVRRGEEAQRRAAGDLTVLIGYDLRYWRELATL
            FGNPYLDFLHRLRVQSVCTVQHLLRRLSELGALWSGHTALVDALARRDPGARALV
            DAYNSHSLALIEGLAGE"
            462..1127
            /gene="SCH10.02"
            462..1127
            /gene="SCH10.02"
            /note="SCH10.02, hypothetical protein, len: 221 aa;
            unknown function, probable CDS suggested by positional
            base preference, GC frame analysis and amino acid
            composition"
            /codon_start=1
            /transl_table=11
            /label="SCH10.02"
            /product="hypothetical protein"
            /protein_id="CAB42011.1"
            /db_xref="PID:e1428992"
            /db_xref="PID:e1428993"
            /db_xref="GI:4753848"
            /translation="MRSPLLACDLMLVPLVDVLTDPNPFARAEALQYDKVLAEGALP
            PVPYQYMPGLSGEAPVAGFDYDALHFLRAHLQVGLPVPYDVLGGDYEQLLLE
            RSTAAQSHLVYHVDHAGAYVPDFPHPLSLDELLAGGPGLSHTLLRELEAVAPAL
            GIDPANFPAPQPPPIAPTELEPAPVAPHDSSPFARERHVMGLHAAATRSIAQSGMI
            VFS"
            complement(1143..1433)
            /gene="SCH10.03c"
            complement(1143..1433)
            /gene="SCH10.03c"
            /note="SCH10.03c, hypothetical protein, len: 96 aa;
            unknown function, probable CDS suggested by positional
            base preference, GC frame analysis and amino acid
            composition"
            /codon_start=1
            /transl_table=11
            /label="SCH10.03c"
            /product="hypothetical protein"
            /protein_id="CAB42012.1"
            /db_xref="PID:e1428993"
            /db_xref="PID:e1428994"
            /db_xref="GI:4753849"
            /translation="MRVLLKATLDEKSNVIRSRMEALIKETVDRIGPRAAYFGTI
            DERRALLVFMQSSQMPPIGEFPFLMELNAEVDLCPIMNGDDLAKGLSLQG"
            complement(1472..2356)
            /gene="SCH10.04c"
            complement(1472..2356)
            /gene="SCH10.04c"

```

```

/note="SCH10.04c, probable RNA polymerase sigma factor,
len: 294 aa; similar to many e.g. IR:005735 (EMBL:U07308),
sigE, Mycobacterium avium extracytoplasmic function
alternative sigma factor (251 aa), fasta scores; Opt: 226
z-score: 266.2 E(): 1.6e-07, 29.8% identity in 168 aa
overlap"
/codon_start=1
/transl_table=11
/label="SCH10.04c"
/product="putative RNA polymerase sigma factor"
/protein_id="CAB42013.1"
/db_xref="PID:e1428994"
/db_xref="PID:g4753850"
/db_xref="GI:4753850"
/translation="MPARSDDMARQFELDRPRLRAVAVRILGSLGADALQEAWLRA
DRANSEVGNFSGWLTIVVARVCLNLLRRARDTRREEPLDDAARQPAATGTAADPAEE
VLADVEGIALIIVLTIDTPAERLAFVLHDFDFDDIAMDRTPAATRLQASRAE
RRVGVPAPEADLPRRRRAVDAYLAATRGDFDALVLIHPDVLSDAAAVVPTPEPV
TVSGMEQVARGAMASMARARSAAVVLVDGMVGMAMENGRRLRVLRLEDFADGRITGI
DVIAPARLNELDITIG"
complement(2472..2672)
/gene="SCH10.05c"
complement(2472..2672)
/gene="SCH10.05c"
/note="SCH10.05c, hypothetical protein, len: 66 aa;
unknown function, probable CDS suggested by positional
base preference and GC frame analysis"
/codon_start=1
/transl_table=11
/label="SCH10.05c"
/product="hypothetical protein"
/protein_id="CAB42014.1"
/db_xref="PID:e1428995"
/db_xref="PID:g4753851"
/db_xref="GI:4753851"
/translation="MTDRYSAAACQGYGGENPEDCGDPVRFVAREHLRAPLRVCPV
HLGSLSLATGVLMPPGIVLR"
complement(2749..3240)
/gene="SCH10.06c"
complement(2749..3240)
/gene="SCH10.06c"
/note="SCH10.06c, hypothetical protein, len: 163 aa;
unknown function, probable CDS suggested by positional
base preference, GC frame analysis and amino acid
composition"
/codon_start=1
/transl_table=11
/label="SCH10.06c"
/product="hypothetical protein"
/protein_id="CAB42015.1"
/db_xref="PID:e1428996"
/db_xref="PID:g4753852"
/db_xref="GI:4753852"
/translation="MRKPTTRTWIAAAVVVVVGGAFYVDGHRVSAAPHAGDPKCD
IVSRLPDDIPGASRDWALGDVAGWQKAVFRCGAELQPNVLCVADGVWDVLDE
ALRRKDGVSVLRTYGRSPAVEFTYSGPREVGGILTALDPAVKWIPQERKCIGLDDAA
TVL"
complement(3412..4695)
/gene="SCH10.07c"
complement(3412..4695)
/gene="SCH10.07c"
/note="SCH10.07c, purA, probable adenylosuccinate

```

```

Query Match      6.6%; Score 110.2; DB 1; Length 39524;
Best Local Similarity 49.8%; Pred. No. 1.1e-07;
Matches 308; Conservative 0; Mismatches 308; Indels 3; Gaps 1;

QY 270 cyagcgtgagagagcggcgccgtctgtccgtgacggcgacggcgccgctcg 329
      ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
Db 33265 CGATTCTGCCGAGATGTCGCCGCCCTGGCGGTGCGGACCGGCGATCACTTCGCCAGG 33324
      ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||
QY 330 tgttccccgccttcgcgcgcgcgtcgagggccgaagacgcgcgtccgcgcgtcgca 389
      ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

```

D	b	33325	GCTTCCCGACACGAGCCCCGGAGAGGTCTCGGAGGCCCGGTGCGGCACACTGGCG	33384
Q	y	390	cggccagtgtaactgtatcgccccgctgcgacctcccgcgcacgaagcgccttag	449
D	b	33385	ACGGCGCGGCAACAGTAGTACCCTCCCGGCGGCTCCCGAGAGTGCGCGCCGCTCG	33444
Q	y	450	cagsgaaccttgtcaagg---cgtgccataagaagtattcggcgcagacgttttctcca	506
D	b	33445	CCGCCACCAACGACGGCGCTACGGCCTGTCTACGACCCCGCACACCGAGTCTGTGTCA	33504
Q	y	507	ccgcgcggaactcagcggatgaagtcaataatccgggtgctggcccagactgccggcg	566
D	b	33505	CCGCCGGCGCCACGAGAGCCATCGCGCGCCCTGCTGCGCCTGCTGAGCCCGCGGACG	33564
Q	y	567	ccaacatactgtctcccgcgcaggttatccaaattacgagcgcgagcgcatccaaca	626
D	b	33565	AGGTCTGTCGCTGGAGCCGTACTACGACTCGTACGCGGCCTGCATCGCATGCGGGCG	33624
Q	y	627	agctggaggtccggcaacttgaacttcattcccgacaaagggtggagatcacatgaat	686
D	b	33625	GCACCCGCGTGCCCGGTGACCTTGGCGCGCAGAGGCGACGTTCCGTCTGACCTGGAGC	33684
Q	y	687	cgtgggaatccatgcgcgaacaagaaccacccgvgatggtcatcataaacccaacaatc	746
D	b	33685	AGCTCGCGCAGCGGTCAACCGACCGACCCGCTGCTGCTGAACACCCCGACAACC	33744
Q	y	747	cgtgcgcgagcggttaactctctacgacatctggccaaggtcgcggaggtggcaaggaagc	806
D	b	33745	CGACGGGCACAGGTCTCTACCCGCGCAGCAGCTGGCCGCGATCGCGAGTGGCGTGGAGC	33804
Q	y	807	tcggaatatgtgatcgctgacaggttttacggcaaatggtttcggcagcgcccgct	866
D	b	33805	CGGACCTGCTGGTGTGACCGACGAGGTCTACGAGCACCTGGTCTCGCACCCGCGAGC	33864
Q	y	867	ttatccgatgggogtctt 885	
D	b	33865	ACATCCCCTGGCGTCTT 33883	
RESULT 10				
AB018112/c				
LOCUS				
DEFINITION				
complete sequence.				
ACCESSION				
NID				
VERSION				
KEYWORDS				
SOURCE				
ORGANISM				
Arabidopsis thaliana				
clone:MAB16.				
Eukaryota; Viridiplantae; Streptophyta; Embryophyta; Tracheophyt;				
euphyllophytes; Spermatophyta; Magnoliophyta; eudicotyledons; co-				
eudicots; Rosidae; eurosids II; Brassicales; Brassicaceae;				
Arabidopsis.				
1 (sites)				
Nakamura,Y.				
TITLE				
Structural Analysis of Arabidopsis thaliana Chromosome 5. IX				
JOURNAL				
unpublished (1998)				
REFERENCE				
2 (bases 1 to 70475)				
Nakamura,Y.				
AUTHORS				
Direct Submission				
TITLE				
Submitted (06-OCT-1998) to the DDBJ/EMBL/GenBank databases.				
Yasukazu Nakamura, Kazusa DNA Research Institute, Laboratory of				
Gene Structure 2; 1532-3, Yana, Kisarazu, Chiba 292, Japan				
(E-mail:yinakam@kazusa.or.jp, Tel:+81-438-52-3935,				
Fax:+81-438-52-3934)				
FEATURES				
Location/Qualifiers				
1..70475				

numbered using the following system eg SC7B7.01c.SC (S. coelicolor), 7B7 (cosmid name), .01 (first CDS), c (complementary strand). The more significant matches with motifs in the PROSITE database are also included but some of these may be fortuitous. The length in codons is given for each CDS.

Usually the highest scoring match found by fasta -o is given for CDS which show significant similarity to other CDS in the database. The position of possible ribosome binding site sequences are given where these have been used to deduce the initiation codon. Gene prediction is based on positional base preference in codons using a specially developed Hidden Markov Model (Krogh et al., Nucleic Acids Research, 22(27):4768-4778(1994)) and the Frameplot program of Bibb et al., Gene 30:157-66(1984) as implemented at <http://www.nih.gov/jp/jun/cgi-bin/frameplot.pl>. CAUTION: We may not have predicted the correct initiation codon. Where possible we choose an initiation codon (atg, gtg, ttg or (att)) which is preceded by an upstream ribosome binding site sequence (optimally 5-13bp before the initiation codon). If this cannot be identified we choose the most upstream initiation codon.

IMPORTANT: This sequence MAY NOT be the entire insert of the sequenced clone. It may be shorter because we only sequence overlapping sections once, or longer, because we arrange for a small overlap between neighbouring submissions. Cosmid 2H4 lies between 2G5 and AH10 on the AseI-B genomic restriction fragment.

Location/Qualifiers  
1..25970  
/organism="Streptomyces coelicolor"  
/strain="A3(2)"  
/db\_xref="taxon:1902"  
/clone="cosmid 2H4"  
1..1822  
/gene="SC2H4.01"  
1..1822  
/gene="SC2H4.01"  
/note="SC2H4.01, probable ATP/GTP binding protein, partial CDS, len: >606 aa; contains PS00017 ATP/GTP-binding site motif A (P-loop)"  
/codon\_start=2  
/transl\_table=11  
/product="putative ATP/GTP binding protein"  
/protein\_id="CAA20595.1"  
/db\_xref="PID:el319219"  
/db\_xref="PID:g3559957"  
/db\_xref="GI:3559957"  
/db\_xref="SPTREMBL:O86584"  
/translation="IDSTAPRAVITSLPTQCTADLCCKGPGGTAGSTTFQPNAA DTDIGYRLTLTSAKAKAVTGKSVTVSDVTPSLGTQVLSVEAKDVRGRWAPAE FSKYAPAGATGTWHDALPGSGVTIAKDTAAEGTRHDATLXTAGAGWTLARRGD ADNSLWLDGSDTTQRTQRAATSAVAVTRDSFTVSAWAYLTDTSQTRVWMAAPCTYG SAFTLYGSAYKKVFNATAGDKVPVLSLGDATAPPLKVVTHLAAYVDTTKDKN KANDTQLFVNGRPGQEPVLDGVSTAVQPVWSSGLOIGRSLVGGAGWGEHFRGLDE VNMQRVLTPELITQAOQLLEGVPANELVAQNDRAASGTITVLSLSPAPSLTSLA AGAVLDENNALVLDGSGASATGVPVVDGTSFVSARVQDLSAKLASKVPVGEALY AGQASAGESSWALWAVKADPVQWKFRTAVAGADGKVTQSAEAPGGDIAETDITWQ VTGVDAQEPWETDPADSAKTETRYGRHLVYGFDDQSETASGFTTPOYGAGALAV GRGRGGTGHLLPCALEELRWVTGAMSADQVRSQVLGGV"  
188..211  
/gene="SC2H4.01"  
/note="PS00017 ATP/GTP-binding site motif A (P-loop)"  
1949..1953  
/note="possible RBS upstream of SC2H4.02"  
1959..8510  
/gene="SC2H4.02"  
1959..8510  
/gene="SC2H4.02"  
/note="SC2H4.02, unknown, len: 2183; very limited similarity to proteins of the Rhs family e.g. WAPA\_BACSU wall-associated protein precursor (2334 aa), fasta scores: opt: 141 z-score: 296.8 E(); 2.7e-09, 23.4% identity in 1237 aa overlap, and RhsA\_ECOLI RhsA protein precursor (1377 aa), fasta scores: opt: 135 z-score: 222.4 E();

FEATURES  
source  
gene  
CDS  
misc\_feature  
RBS  
gene  
CDS

3.8e-05, 23.0% identity in 1016 aa overlap. Contains possible membrane anchor around aa 1900. Alternative start at aa 27 would give N-terminal signal sequence"  
/codon\_start=1  
/transl\_table=11  
/product="hypothetical protein SC2H4.02"  
/protein\_id="CAA20596.1"  
/db\_xref="PID:el319220"  
/db\_xref="PID:g3559958"  
/db\_xref="GI:3559958"  
/db\_xref="SPTREMBL:O86585"  
/translation="MRHPLGVPGTGPGRHRRRAVRPMRRVSAVVGIALLPGLLA PAATADDVPLGRPELSTPRAOHVAPFKARPDAATAVVRDSARQASAGRAVKDRS RGTSTAGSARLTASGADEAEPGRPLVLAAPAAKAKRADRVKVDVLDQKAARP LGVGVILKVTGPADGAARJGIDISAFAYAGDWAAGRLQVLRDCTDACPAPPV AFTNBERGERTLADIAFAKPAAGSTMTLALAAAGTASADGDKATPLAPSSWAGGS SGSTFWSPILKAPAPAGEAKLEISYDGSVDGRTSTNNQGTSGVGGFTISFIE RYGSCTDDHADRHDLCKYDNALSKGATELVKDDTQGWRLKNDKADSKVTHST GADNGDDNGEYVTTTGGTFRVFGONKLDGAAADVTKSVWTVFVGDDEGPGYAD GTGFTSRDEKQAWRNLLYAEETHGNATYVWEDTNNYDKLGGDDTGTSYRGVYLK EIRGQRAGALFSFGTPQASHRILVLDGERCLASGTGCDALTEDTNDNMPDVPFRICK DDCKPYLDSPFFTRKMLTISVSYVAMDAATPAFSPVDTWTLHLXLDPGDGTDS DQSLWDEIKHTGRGTDLSDAVKSFVMPNRVDPGSDIILDFERPLRVSVETG AQTIVDYLPAFCVAGQIMPKADENTKCFPVWSPYGOEPIIDWFQYPPVSSVTTD PLGSGAVQHTYEXSGGAWHYNDPLTPAKERTWSQWRGKVTHTYTPSGSTRAKT VTYLGRMGDRVLGGCKTDPDRRRKAEVGVTAATAVSDOLAGFQRESVADGD KEVSGTVNDPMSKRTATOKHSYADTEAYYRVGASHARTIISRNPDRVTRITGY DDYGNPVEDLGGDSDVTGDEKTNIAFARNDAGLTALVSRIVGRACSVTSDALD LPADARPGDVVSDKATADTTTSATQKPTKGDAWGRAGKYGDDRPLMQTTAVT DYTILREVQVRNTDDVTISKTETPPAAGPLTSTVYNAKHRTTIVKDFALGADLK VTDANGKTESAYDSLGVTSWLPNRSALGKTANYVYGVSKVTSPLPWSSATLNG DGSVYRTYEVISLRTROYQAPSAQGRVIAOTLYDGRGLPFTAOADIMDDTAAPA GKIVLDGGQAPROTDSYDGMGRVTAKYSYGVTEWADYTAGDLVLGTGAPGGGS ANAVTDFAGTRVRRDITAGTQAGTDYMTIRYAFDAADROKSIHAHRSAWTYTDL FGRQVSTDPDKGTTVEYDALDRAVSTDCRGEVLLFEYDLGRTGMQWSAKTGAN KLAAMSFDTLAKGQQTAVRYEGGETGRATYQKVTYDPLKYKTNELTLPANDPLVA AGVPLADLQSTGYNLDTGVKQAAAPAVAGLSAETVSYTYDGLGVLTAKTGTLYQQA AYSPLGDFURQMTLATDPTGAKVYLNDYEAGTRELTRSYVTDVHGMLOELKQQD DAGNITSVSDATILGGTGKADHQCTYDGRHRLSEAWTPETADCSSTRTVAGLGAA PYWTSYQDSSGLRSKQFEHRMSGDDVTYEYGAEGPHALSATVGAENASTYTD TDGNTETPGVRAQTLDWNAEGRLAGVSEPAAGKPKATYAVYVADAGDILLIRPTT TDGETVLYLTTEVHLKVSGNAKALSGARTYKAGSAVIAVTRTAGVSGTKLFLA GDHRTSGLAINDTAFKARWSTPFGAPRGTAGMDPDRLGKPADATGLTGLG AYOQDPDTGRLSVDPDLEPKPNTLNGYAVASNSPTNSDPSCGSDGLGLGALGA IIGVGVGAVIGAGAAITAVGSLGGGGGGGCTAPTSSGGWTQPLTKWTPGATYNF ITKSWDLPFPNSQSLSEMLASMPDWGIVDPKAANRWETSRSLFFLWGGGYPFRE HQDFRGGDAFTSILAQDDTISGLRSKMGVQARKGTGFAKEVGFYVDEGPEGSPW YKFNLSRGAADIAVLNTGAVGTGNQADAFLGTSYSGKARISINKEGSGVTLKFSAW NGSDWRGATHVVRPNWPAFEDTFGAAVREDFSWEEKWPINECVNYSWLE"  
8536..8571  
/note="hairpin loop with 17/17 bp stem"  
8602..9063  
/gene="SC2H4.03"  
8602..9063  
/gene="SC2H4.03"  
/note="SC2H4.03, possible lipoprotein, len: 153 aa; contains possible N-terminal signal sequence with appropriately positioned PS00013 prokaryotic membrane lipoprotein lipid attachment site"  
/codon\_start=1  
/transl\_table=11  
/product="putative lipoprotein"  
/protein\_id="CAA20597.1"  
/db\_xref="PID:el319221"  
/db\_xref="PID:g3559959"  
/db\_xref="GI:3559959"  
/db\_xref="SPTREMBL:O86586"  
/translation="MLKTPFAPAVRWTAALVITLLAATACDDTGARAATVPADRVTGA WQAGSGALTILSADRTTASGLDSKGLGCTPGDGTGGDAFWAFADDGDLMYSPRA RSGSEIGLSRAEQSDPECRULTVVDGDTLCAITSDPETPCDLGVRRTRKD"  
8647..8679  
/gene="SC2H4.03"  
/note="PS00013 Prokaryotic membrane lipoprotein lipid

[illegible]

RESULT .12

SAAJ5198/c	3240 bp	DNA	BCT	04-SEP-1998
LOCUS				
DEFINITION	Streptomyces alboniger stgA, stgU and stgR genes.			
ACCESSION	AJ005198			
NID	g3550621			
VERSION	AJ005198.1	GI:3550621		
KEYWORDS	LysR-type transcriptional regulator; stgA gene; stgR gene; stgU gene.			
SOURCE	Streptomyces anulatus.			
ORGANISM	Streptomyces anulatus			
REFERENCE	Bacteria; Firmicutes; Actinobacteria; Actinobacteridae; Actinomycetales; Streptomycineae; Streptomycetaceae; Streptomyces.			
AUTHORS	Jimenez, A.			
TITLE	Direct Submission			
JOURNAL	Submitted (04-JUN-1998) Jimenez A., Universidad Autonoma de Madrid, Facultad de Ciencias, Centro de Biologia Molecular 'Severo Ochoa' (CSIC/UAM), Cantoblanco. Madrid, 28049, SPAIN			
REFERENCE	2 (bases 1 to 3240)			
AUTHORS	Tercero, J.A., Espinosa, J.C. and Jimenez, A.			
TITLE	StgR, a new Streptomyces alboniger member of the LysR family of transcriptional regulators			
JOURNAL	Unpublished			
FEATURES	Location/Qualifiers			
source	1..3240			
	/organism="Streptomyces anulatus"			
	/db_xref="taxon:1892"			
	complement(180..1388)			
gene	/gene="stgA"			
CDS	complement(180..1388)			
	/gene="stgA"			
	/function="putative aminotransferase"			
	/codon_start=1			
	/transl_table=11			
	/product="StgA protein"			
	/protein_id="CAA06435.1"			
	/db_xref="PID:ej318407"			
	/db_xref="PID:g3550622"			
	/db_xref="GI:3550622"			
	/db_xref="SPTREMBL:O88124"			
	/translation="MEPRQSKLSEVCYELRGPVIEHANALEPAGHSVLRNTGNPAL FGPEAPEIVQDMIRMLPOAHGYTDSRGVLSARRAORYCALGLDEVSDDVFLGNDM VSELVMAVALLEDGDEIILPAPDFLWATVTLSCGKAVHYVCDAAWYDPDLDDM ASKITDQTAIEIVLITGAVYKPEILLEGILDLARHGLAWFADEIYDQILYDDRVHHS VAALPDLVLTITCGSLTKYRAGFSRWGLVITGPQROHARDYLEGLTMLASMLCPNNA PAQFAIQALGSGQSIRELPAGCALHEQRDRAWEKLEIPGVCVKPKGALYAFAPRL DPKVHPITHDKEKFLVLLQEKIQVVGQTFNPNRPDPHFRILTLPHADDDLAAISRIG RFLSGYRQGR"			
gene	1473..2105			
	/gene="stgU"			
CDS	1473..2105			
	/gene="stgU"			
	/codon_start=1			
	/transl_table=11			
	/product="StgU protein"			
	/protein_id="CAA06436.1"			
	/db_xref="PID:ej318408"			
	/db_xref="PID:g3550623"			
	/db_xref="GI:3550623"			
	/db_xref="SPTREMBL:O88133"			
	/translation="MSPRRSYDQYCAARALDADVGDRWTLTVVRELIVGPPRYTDLH DLPGVSDVLASRLKMDPEQGLTFRRLPPAAAYVVELTLKROLLPVLEALAAWG HGLSERPTDAVRHFAWFLPMLFTLRIGAECTVEVGLGGRFHVRLGDGAETYGDG APDAPEVRLAUDACTCVAVSRGELTVGEGVRAGRIELSGEGTLAKALLA"			
gene	2219..3202			
	/gene="stgR"			
CDS	2219..3202			
	/gene="stgR"			
	/note="LysR-type transcriptional regulator"			
	/codon_start=1			
	/transl_table=11			
	/product="stgR protein"			
	/protein_id="CAA06437.1"			

[illegible]

Query Match 6.3%; Score 104.8; DB 1; Length 3240;  
Best Local Similarity 51.4%; Pred. No. 8.4e-07;  
Matches 300; Conservative 0; Mismatches 27; Indels 12; Gaps 2;

QY 395 cagttcaagtctacgcgcccggcggtcgcctcccccgcgcgaagcgcgttagcacag 454  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 1202 CAGGCGACGGCTACACGGACTCGCCGGCGGTCTTCCGCCCGCGGCCTGCCCGAG 1143  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
QY 455 cacttgtcacaggcgttcacctacaagctatggccgcagcagcttctctaccgccg 514  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 1142 CGTATCAGCGCGTCGCCCTCGACGAGGCTCTCGGTGAGAGCTTCTCTCGCAACGCGC 1083  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
QY 515 ggaactcagcgcatcagaagtcaataatcccggtgctgccccagcagtcgcccgaacata 574  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 1082 GTGTCGAGTGCTCAGCATGGCGGTCCAGGCGCT---CCTGGAGAGCGCGCACAAATC 1026  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
QY 575 ctgcttccccgcgcaagctatcaaatcagagggcgagcggcattcaaacagtgag 634  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 1025 CTATCCCCCGCGCCGACATCCCGCTGTGGACGGGGTGACGACCCTCTCCGGAGGCAAG 966  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
QY 635 gtccggcacttcgactcatcccccgaaggggtggtgagatgcacatgcgtctgaa 694  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 965 CCCTCCACTACGCTCTCGACGAGGCGGCGACTGTATTCCCGACTCGACGATGCCC 906  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
QY 695 tccatcgcgcagaacacacccgcgctggttcataataaacccaacaatccgtgcgcg 754  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 905 TCGAAGATCACCGACACGAGCGCGTCGAGATCATCAACCCCA-----CGGCGC 855  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
QY 755 agcgttactctacacacatctggcccaaggtgcgaggtggcgaagctcggaata 814  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 854 CGCGTCTATCCGAAGAGATCTCTGGAGGACATCTCTGATCTCGCGCGGCGACGCGCTG 795  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
QY 815 ttggtgatcgtacagagttaacgcaaacgttgttctggcagcgcctgttatcccg 874  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 794 ATGGTTCTTCCCGACGAGATCTACGACCAAGTCTCTAGCAGACGCGCTGCACACAGC 735  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
QY 875 atggcggtctttggcacatggcccggtctgttcattggtatctctccaaagctgtgg 934  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 734 GTGGCGCCTCGCCCGGATCTCGTGGTCCCTCACCTCTCGGGTCTTTCGAGACGTAC 675  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
QY 935 atagtgcgtgattggcgaactgattgggtggcgggtgtacgacc 978  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||  
Db 674 CGCGTGGCGGTTTTCGCTTGGCTGGCTGGCTGGCTGGCTGGCTGGCTGGCTGGCTGGCT 631  
||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| ||||| |||||

RESULT 14  
MSGBI970CS/c  
LOCUS MSGBI970CS 39399 bp DNA BCT 26-JUN-1996  
DEFINITION Mycobacterium leprae cosmid B1970 DNA sequence.  
ACCESSION L78815  
NID G137775  
VERSION L78815.1 GI:137775  
KEYWORDS  
SOURCE Mycobacterium leprae (clone: cosmid B1970) (tissue library: Lorist 6) DNA.  
ORGANISM Mycobacterium leprae  
Bacteria; Firmicutes; Actinobacteria; Actinobacteridae;  
Actinomycetales; Corynebacterineae; Mycobacteriaceae;  
Mycobacterium.  
REFERENCE  
1 (sites)  
Eiglmeler K., Honore N., Woods S.A., Caudron B. and Cole S.T.  
Use of an ordered cosmid library to deduce the genomic organization  
of Mycobacterium leprae  
Mol. Microbiol. 7 (2), 197-206 (1993)  
REFERENCE  
MEDLINE 93188700  
AUTHORS Smith D.R., Richter P., Rubenfield M., Butler C., Lee H.-M., Xu O., Gunderson K., Chung M., Maher J.K., Deloughery C., Aldrich T., Imrich J., Tulig C., Smyth A., Drill S., Avrukh A.S., Rice P., Abendschan K., McDougall R., Deloughery C., Kirst S., Safer H., Connelly S., McDougall S., Eiglmeler K., Bergh S., Cole S., Robinson K., Jaehn L., Gryan G., Church G.M. and Mao J.  
Prepublication submission  
TITLE





gene	9624..10881	/gene=F9D16.20"	
CDS	join(9624..9967,10214..10330,10428..10881)		
	/gene=F9D16.20"		
	/note="similarity to AR411 - Arabidopsis thaliana		
	PATX:G1669603"		
	/codon_start=1		
	/product="putative protein"		
	/protein_id="CAA23021.1"		
	/db_xref="PID:el1386934"		
	/db_xref="PID:G4454024"		
	/db_xref="GI:4454024"		
	/translation="MDGELEATVIRVYSGSDAFSGESSCTFSPSFCLPMETSSFFVEP		
	EMETSGDELEGELKPYRPFSTQVTLISSVSLPDSKPFRRDDKKQBSHGCLLSNGSRA		
	DHIRISSEKSKKKKQKRVQEVKEENLLSDAWKRYGQKPIKSGPSYRISYGRS		
	SSKGLCARQVKNPQRFETTYTNEHHHELPTNRNSLAGSTRAKTSQPKPTLTK		
	SEKIVSPKTSNPMPISADSSSVAVQMSVAETSTHQAGAIEGRRLSNGLSLDLMSG		
	SGTSPSTGDFDELLNSQEFFSGYLWNY"		
intron	9968..10213	/gene=F9D16.20"	
	/number=1		
exon	10214..10330	/gene=F9D16.20"	
	/number=2		
intron	10331..10427	/gene=F9D16.20"	
	/number=2		
exon	10428..10881	/gene=F9D16.20"	
	/number=3		
repeat_region	10899..10931	/note="33bp ta tandem repeat"	
exon	complement(11426..11752)	/gene=F9D16.30"	
	/number=1		
gene	11426..13581	/gene=F9D16.30"	
CDS	complement(join(11426..11752,11828..11994,12098..12206,		
	12305..12457,12570..12647,12732..12932,13067..13279,		
	13390..13581))		
	/gene=F9D16.30"		
	/note="strong similarity to cellulase - Phaseolus		
	vulgaris, PID:G1039431; contains EST gb:F14141, T76139,		
	F13971"		
	/codon_start=1		
	/product="putative cellulase"		
	/protein_id="CAA23022.1"		
	/db_xref="PID:el1386935"		
	/db_xref="PID:G4454025"		
	/db_xref="GI:4454025"		
	/translation="MGKLLVLMYGMFLAFESLEALEYGDALNKSLIFFEQSGKGLP		
	TNQVRKWRADSGALSDGLANVLIGGYDAGDNKVFYPMKSFPTLLSWAAIEYQNEI		
	SNVNLQGLRSTIKWGTILRAHTSPNMLYQVGDGNSHSCWPEREDMDTISRLYS		
	ISSSPGSEAGEAAGAAASLVKFSVDSTSLTNHAKTLFEFADKRGYSQASC		
	PFFYCSYQDELLAAAWLYKAGDKIYINVISNKDWQSOAVNEFSWDNKFYGAQAL		
	TVLFHYSKTLKAGVSIQCGSTFTVSIQINFASKQVDYILGNPNMKMTYWGFGTK		
	YPTOPHRRGSSLPSTQSKPEKIDCGNGYSYYSNSTPNPNVHIGAIVGPNSSDQYSDK		
	KDSYSHAEPYVINAFTGPPVAALISSG"		
	complement(11753..11827)		
intron	/gene=F9D16.30"		
	/number=1		
exon	complement(11828..11994)		
	/gene=F9D16.30"		
	/number=2		

Query Match 5.7%; Score 94.2; DB 7; Length 119430;  
Best Local Similarity 56.5%; Pred. NO. 1.9e-05;  
Matches 196; Conservative 0; Mismatches 148; Indels 3; Gaps 1;  
Qy 440 aggcgcgtagcagagcactgttcacagggcgtgcctacaagctatcggcgcagcactc 499  
DB 29109 AGTGGCGTGGACAGACTATCTAAACAAGCTCTTCCAAAGAACTTACGCGACATGACGCTG 29168

Qy	500	ttctcaaccgcccgggaactcagcgcatcgaatcataatcccggtgctggcccagact	559
Db	29169	TTTATGACTCTGGGATGCAACAAGCTATTAGCTCGGGTAGACATTCTCGCTAACC	29228
Qy	560	gcggcgcccaactactgcttccccgcgagcgatccaattacgagcgcgagcgga	619
Db	29229	AAAG--CCAACGTTTTCCTCCGAGTCCGGCTTCCATGGGACCTAGTCCGCTCCATC	29285
Qy	620	ttcaacaagctggaggctccggcaattcgacctatccccgaagaagggtgggaatcgac	679
Db	29286	TACAAGAACCTTGAGTCCGGCCACTATAATTTCCTCCAGAAAAAAGACTTTGAATCGAC	29345
Qy	680	atcgactcgctggaaatccatctcgcgcaagaaccacccgcatgggtcatcataaaccca	739
Db	29346	TTTGATAGGCTCCGAGGCTCGTGGAGAGAACACATTTCGCAATTATTATCAACCCC	29405
Qy	740	aacaatcgtgcggcagcggtttactcttaecaccatctggccaaggt	786
Db	29406	CACAAGCCCAATGGTACACACCTACTCCGAGGCTCATCTCAAACAGGT	29452

Search completed: October 1, 1999, 15:32:18  
Job time: 6041 sec

---

